

# Calculation of Nucleon Spin in Lattice QCD





**Takumi Doi**

(Nishina Center, RIKEN)

$\chi$ QCD Collaboration

M. Deka, Y.-B. Yang, B. Chakraborty, S.-J. Dong, T. Draper,  
M. Glatzmaier, M. Gong, H.-W. Lin, K.-F. Liu, D. Mankame,  
N. Mathur, T. Streuer

[arXiv:1312.4816](https://arxiv.org/abs/1312.4816)



- Outline

- Introduction
- Lattice QCD framework
  - Brief review of Lattice QCD
  - Nucleon spin on the Lattice
- Lattice QCD results
- Summary & Prospects

# Nucleon structure from QCD

- Nucleon: the only hadron which is stable
  - the structure is crucial to understand nucleon itself, QCD, (& beyond SM)
  - ***Electric/Magnetic structure***


$$\left(\frac{d\sigma}{d\Omega}\right) = \left(\frac{d\sigma}{d\Omega}\right)_{Mott} \left[ \frac{G_E^2 + \tau G_M^2}{1 + \tau} + 2\tau G_M^2 \tan^2 \frac{\theta}{2} \right]$$

- $G_E$ : electric form factor
- $G_M$ : magnetic form factor

- ***Deep Inelastic Scattering (DIS)***

$$\left(\frac{d^2\sigma}{d\Omega dE'}\right) = \left(\frac{d\sigma}{d\Omega}\right)_{Mott} \left[ W_2 + 2W_1 \tan^2 \frac{\theta}{2} \right]$$

- $W_1, W_2 \rightarrow F_1, F_2$  structure functions



# Puzzles in Nucleon structure

- Do we know precisely ? Flavor/Glue DoF ?

- **Vector form factor**

- One of the most well-determined quantities, but... → “proton size crisis”

- Strangeness element  $\leftrightarrow$  constrain  $G_A^s$



(2010/07)

- **Scalar form factor**

- Origin of the mass

- pi-N-Sigma term  $\leftrightarrow$  pi-N int., rho mass shift in medium

- Strangeness element  $\leftrightarrow$  Dark Matter Search

- **More for Beyond SM**

- EDM form factor  $\leftrightarrow$  (strong) CP problem

- Tensor form factor  $\leftrightarrow$  Non V-A Int in beta-decay

# Puzzles in Nucleon structure


- Spin (axial vector)

– “Spin crisis”

- quark spin is small !

$$\Delta\Sigma = \sum_q [\Delta q + \Delta\bar{q}] = 0.2-0.3$$



EMC(1988)



$$g_1(x) \simeq \frac{1}{2} \sum_q e_q^2 [\Delta q(x) + \Delta\bar{q}(x)]$$

– Glue ?





$$\int_{0.05}^{0.2} \Delta g(x) dx = 0.1 \pm_{0.07}^{0.06}$$



# Where does the proton spin come from ?

$$\frac{1}{2} = \frac{1}{2} \Delta\Sigma + L_q + \Delta G + L_G$$

- **Quark spin: 20-30%**
  - DIS, Lattice
- **Glue spin: ~20% ?**
- **Quark orbital angular momentum:**
  - Small in Lattice ? (for a part of diagrams)
- **Glue orbital angular momentum ?**
  - **→ Dark-Spin ?**



- Outline

- Introduction

- **Lattice QCD framework**

- Brief review of Lattice QCD

- Nucleon spin on the Lattice

- Lattice QCD results

- Summary & Prospects

# QCD (DoF=quarks/gluons)

- Formula of QCD: very simple & beautiful

$$\mathcal{L} = -\frac{1}{4}G_{\mu\nu}^a G_a^{\mu\nu} + \bar{q} [\gamma^\mu (i\partial_\mu - gA_\mu) - m] q$$

$$G_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + g f_{abc} A_\mu^b A_\nu^c$$


- Only 4 parameters      quark masses ( $m_u, m_d, m_s$ )  
coupling constant  $\alpha_s = g^2/4\pi$

mass ( $\overline{MS}, \mu = 2\text{GeV}$ )	$m_u$	$m_d$	$m_s$
[MeV]	$2.3^{+0.7}_{-0.5}$	$4.8^{+0.5}_{-0.3}$	$95 \pm 5$

(PDG2013)

- Solving QCD: very challenging

- Coupling is “strong” at low energy
- Nonperturbative effects
- Quantum effects w/ infinite # of DoF




For quark mass determination by lat, see, e.g.,  
T.Blum et al. (RBC Coll.) PRD82(2010)094508

# Lattice QCD

## First-principles calculation of QCD

$$Z = \int dU dqd\bar{q} e^{-S_E}$$



quarks on sites  
 $q(x)$

gluons on links

$$U_\mu(x, x + \mu) = \exp[-iaA_\mu]$$




K.G. Wilson

- Well-defined regularized system (finite  $a$  and  $L$ )
- Gauge-invariance manifest
- Fully-Nonperturbative
- DoF  $\sim 10^9 \rightarrow$  Monte-Carlo w/ Euclid time

# Lattice QCD

## First-principles calculation of QCD

$$Z = \int dU dq d\bar{q} e^{-S_E}$$



quarks on sites  
 $q(x)$

gluons on links

$$U_\mu(x, x + \mu) = \exp[-iaA_\mu]$$




K.G. Wilson

- (1) Generate the QCD vacuum (configurations  $\{U\}$ ) with the appropriate Monte-Carlo weight ( $\exp(-S_E)$ )
- (2) Measure the observable of your interest

$$\langle \mathcal{O} \rangle = \frac{1}{Z} \int dU dq d\bar{q} \mathcal{O} e^{-S_E} = \frac{\# \text{configs}}{\sum_i} \mathcal{O}[U^i]$$

The QCD Vacuum” is highly non-trivial and non-perturbative


# The QCD Vacuum” is highly non-trivial and non-perturbative



By D.B. Leinweber

<http://www.physics.adelaide.edu.au/theory/staff/leinweber/VisualQCD/Nobel/index.html>

# Measurement on the QCD vacuum



Solve propagator


$$D[U](z, x) D^{-1}(x, y) = \delta_{z, y}$$

Calc correlator

$$\begin{aligned} & \langle 0 | N(t) \bar{N}(0) | 0 \rangle \\ &= \sum_n \langle 0 | N(t) | n \rangle \langle n | \bar{N}(0) | 0 \rangle \\ &= \sum_n |A_n|^2 \exp(-E_n t) \quad (A_n = \langle 0 | N | n \rangle) \\ &\rightarrow |A_0|^2 \exp(-E_0 t) \quad (t \rightarrow \infty) \end{aligned}$$

# Significant advances in Lattice QCD


## Software (theory)



New algorithms make physical point calc possible !

Estimate @ Lat2001 at Berlin  
"Berlin Wall"



## Hardware



K-computer  
(10PFlops)

# Status of Lattice QCD

*Hadron spectrum well reproduced !*




**Fully dynamical QCD simulations  
at the physical quark mass point  
already performed**

PACS-CS Coll., PRD81(2010)074503  
BMW Coll., JHEP1108(2011)148

## Next Challenge




## Another direction (HAL Coll.)



## Nucleon Dirac Radius

$$\langle r_1^2 \rangle$$

$$F_1^{n-d}(Q^2) \approx F(0) \left[ 1 - \frac{1}{6} Q^2 \langle r_1^2 \rangle + \mathcal{O}(Q^4) \right]$$



ChPT predicts divergence  $\sim \log m_\pi^2$

Larger  $L_g$ , smaller  $Q_{\min}^2$  are desirable

## Isovector matrix elements


review talk by S.Syritsyn @ Lat13

## Drama of the Axial Charge

$$g_A$$

$$\langle N(p) | g \gamma^\mu \gamma^5 q | N(p) \rangle = g_A u_p \gamma^\mu \gamma^5 u_p,$$


Experiment (W.A.) [PDG'12]  $g_A^{\text{ave}} = 1.2701(25)$



Many lattice calculations underestimated  $g_A$  by 10-15%


## Strangeness EM form factor

## Magnetic




$$G_M^S(0) = -0.017(25)(07)$$

## Electric



TD et al., PRD80(2009)094503

# How about proton spin ?



( $\overline{\text{MS}}$ ,  $\mu=2\text{GeV}$ )

Fig from C. Alexandrou et al., PRD88(2013)014509

# Formulation on the Lattice

- 1st-moment  $\langle x \rangle$  and spin  $J$  studied simultaneously
- Matrix elements of **energy-momentum tensor**

*Gauge invariant decomposition*

X.Ji (1997)

$$T_q^{\mu\nu} = \frac{i}{4} [\bar{q} \gamma^\mu \vec{D}^\nu q - \bar{q} \gamma^\mu \overleftarrow{D}^\nu q + (\mu \leftrightarrow \nu)]$$

$$T_g^{\mu\nu} = \frac{1}{4} g^{\mu\nu} F^2 - F^{\mu\alpha} F^\nu{}_\alpha$$

$$T_q^{\mu\nu} \rightarrow \bar{q} \vec{\gamma} \gamma_5 q + \bar{q} [\vec{x} \times (-i\vec{D})] q$$

$$T_g^{\mu\nu} \rightarrow \vec{x} \times (\vec{E} \times \vec{B})$$

**Recent developments:**

Chen et al., Wakamatsu, Hatta, Leader & Lorce, ...

- Nucleon matrix elements

$$\langle p, s | T^{\mu\nu} | p', s' \rangle = \bar{u}(p, s) \left[ T_1(q^2) \gamma^\mu \bar{p}^\nu + T_2(q^2) \bar{p}^\mu i \sigma^{\nu\alpha} / 2m \right. \\ \left. + T_3(q^2) (q^\mu q^\nu - g^{\mu\nu} q^2) / 2m + T_4(q^2) g^{\mu\nu} m / 2 \right] u(p', s')$$

$$\langle x \rangle = T_1(0)$$

$$J = \frac{1}{2} [T_1(0) + T_2(0)]$$


(angular) momentum sum rules

$$\langle x \rangle_q + \langle x \rangle_G = 1 \quad J_q + J_G = 1/2$$

# Formulation on the Lattice

- Calculate 3pt (& 2pt) → matrix elements

$$\begin{aligned} \Pi^{3pt}(\vec{p}, t_2; \vec{q}, t_1) &= \sum_{\vec{x}_2, \vec{x}_1} e^{-i\vec{p}\cdot\vec{x}_2} e^{+i\vec{q}\cdot\vec{x}_1} \langle 0 | \mathcal{T} [J_N(\vec{x}_2, t_2) T^{\mu\nu}(\vec{x}_1, t_1) \bar{J}_N(0)] | 0 \rangle \\ &\quad (T^{\mu\nu} = T^{4i}) \end{aligned}$$







- Typical examples:

$$\text{Tr} \left[ \Gamma_e \Pi_{T_{4i}}^{3pt}(\vec{p}, t_2; \vec{q} = \vec{0}, t_1) \right] = C \cdot e^{-m(t_2-t_1)} e^{-Et_1} [2p_i \cdot T_1(0)]$$

$$\begin{aligned} \text{Tr} \left[ \Gamma_m \Pi_{T_{4i}}^{3pt}(\vec{p} = \vec{0}, t_2; \vec{q}, t_1) \right] \\ = C \cdot e^{-m(t_2-t_1)} e^{-Et_1} \left[ -i\epsilon_{ijm} q_j (T_1(-q^2) + T_2(-q^2)) \right] \end{aligned}$$

- Other momentum combinations are calculated and  $T_1, T_2, (T_3)$  are determined simultaneously



- Outline

- Introduction

- Lattice QCD framework

- Brief review of Lattice QCD

- **Nucleon spin on the Lattice**

- Challenges: Disconnected Insertion (DI) and Glue


- Lattice QCD results

- Summary & Prospects


# Challenges in Lattice QCD

## (1) Disconnected Insertion (DI)

- Two kinds of calc in Lattice:




Connected Insertion (CI)




Disconnected Insertion (DI)

- DI is inevitable for flavor singlet quantities, but...
  - **All(source)-to-all(sink)** propagator is necessary
  - Straightforward calculation **impossible**
    - **$O(10^9)$  inversions** for  $O(10^9) \times O(10^9)$  matrix

$$\text{Tr}[\Gamma M^{-1}] = \boxed{\sum_x} \text{Tr}_{\text{color}}^{\text{spin}} [\Gamma M^{-1}(\underline{x}, \underline{x})]$$



# How about proton spin ?



( $\overline{\text{MS}}$ ,  $\mu=2\text{GeV}$ )

Fig from C. Alexandrou et al., PRD88(2013)014509

# The approach for disconnected insertion

- **Stochastic Method for DI**

- Use  $Z(4)$  (or  $Z(N)$ ) noises such that

$$\lim_{L \rightarrow \infty} \frac{1}{L} \sum_{l=1}^L \eta_i^{l\dagger} \eta_j^l = \delta_{ij}$$

- DI loop can be calculated as


$$\text{Tr}[\Gamma M^{-1}] = \lim_{L \rightarrow \infty} \frac{1}{L} \sum_{l=1}^L \eta^{l\dagger} (\Gamma M^{-1} \eta^l)$$

S.-J.Dong, K.-F.Liu,  
PLB328(1994)130

- Introduce new source for noises (“off-diagonal” part)

- $\rightarrow$  **Unbiased subtraction** using **hopping parameter expansion (HPE)**
- Off-diagonal contaminations are estimated in unbiased way

c.f. other approaches  
All-to-all (Foley et al., 2005)  
CAA/AMA (Blum et al., 2012)



# Stochastic method for DI

- Stochastic Method for DI




S.-J.Dong, K.-F.Liu,  
PLB328(1994)130

- Noise


$$\lim_{L \rightarrow \infty} \frac{1}{L} \sum_{l=1}^L \eta_i^{l\dagger} \eta_j^l = \delta_{ij}$$

- DI loop

$$\text{Tr}[\Gamma M^{-1}] = \lim_{L \rightarrow \infty} \frac{1}{L} \sum_{l=1}^L \eta^{l\dagger} (\Gamma M^{-1} \eta^l)$$



+




**Stochastic source**

**Signal part**

**Noise part**

# Improvement of DI calc

- The unbiased subtraction using hopping parameter expansion (HPE) to eliminate off-diagonal noises




→ The error reduces by a factor of 2 or more

# Challenges in Lattice QCD

## (2) gluon matrix elements

- Gluon operator

$$T_G^{\mu\nu} = \frac{1}{4} g^{\mu\nu} F^2 - F^{\mu\alpha} F^\nu{}_\alpha$$



– Implementation is simple w/ link variables

$F_{\mu\nu} \leftrightarrow$  clover term w/ link  $U_\mu$

– In practice, **S/N is known to be notoriously noisy**

- Gluon DoF fluctuate too much in high-freq mode

M. Gockeler et al.,  
Nucl.Phys.Proc.Suppl.53(1997)324

# The approach for Glue

- Field tensor constructed from overlap operator

$$F_{\mu\nu}(x) \longleftarrow \text{Tr}_{(\text{spinor})} [\sigma_{\mu\nu} D_{ov}(x, x)]$$

( $a \rightarrow 0$ )

K.-F.Liu, A.Alexandru, I.Horvath  
PLB659(2008)773

$$D_{ov} = \rho \left( 1 + X \frac{1}{\sqrt{X^\dagger X}} \right), \quad X = -\rho + D_W$$

- Ultraviolet fluctuation is expected to be suppressed (automatic smearing)
- In order to estimate  $D_{ov}(x, x)$ , stochastic method is used w/ color/spinor & (some) spacial dilution





$$D_{ov}(x, x) \Leftarrow \langle \eta_x^\dagger (D_{ov} \eta)_x \rangle$$

c.f. other approaches

Smearing (Meyer et al., 2008)

Change Action & response (Horsley et al., 2012)

Wilson-Flow (H.Suzuki, 2013)



- Outline

- Introduction
- Lattice QCD framework
  - Brief review of Lattice QCD
  - Nucleon spin on the Lattice
- **Lattice QCD results**
- Summary & Prospects

# Lattice Setup

- **Wilson Fermion** + Wilson gauge Action
  - 500 configs with **Quenched approximation**
  - $1/a=1.74\text{GeV}$ ,  $a=0.11\text{fm}$  ( $\beta=6.0$ )
  - $16^3 \times 24$  lattice,  $L=1.76\text{fm}$
  - $\kappa(\text{ud}) = 0.154, 0.155, 0.1555$ 
    - $m(\pi) = 0.48, 0.54, 0.65 \text{ GeV}$
    - $m(\text{N}) = 1.09, 1.16, 1.29 \text{ GeV}$
    - $\kappa(\text{s})=0.154$  ,  $\kappa(\text{critical})=0.1568$


# Lattice Setup (cont'd)

- Disconnected Insertion (DI)
  - Z(4) stochastic method, #noise=500
  - Unbiased subtraction w/ up to 4th HPE
- Glue matrix element
  - Overlap operator  $D_{\text{ov}}(x,x)$
  - Z(4) stochastic method, #noise=2, w/ color/spinor dilution + spacial dilution (d=2 & even/odd  $\rightarrow$  taxi-distance=4)
- Improvement
  - Many nucleon sources, #src=16
  - CH, H and parity symmetry:
    - $(3\text{pt})=(2\text{pt}) \times (\text{loop}) \rightarrow (3\text{pt}) = \text{Im}(2\text{pt}) \times \text{Re}(\text{loop}) + \text{Re}(2\text{pt}) \times \text{Im}(\text{loop})$

# Results for CI: $q^2$ -dependence


$$T_1 + T_2$$

$[T_1+T_2](q^2)$  vs  $T_1(q^2)+T_2(q^2)$  for up and down quark (CI)



$$T_1(u+d), T_2(u+d)$$

$T_{1,2}^u(q^2) + T_{1,2}^d(q^2)$  for Connected Insertion



From dipole-fit

From forward matrix element

Dipole-fit performed

$m_\pi = 0.48\text{GeV}$

Different  $q^2$  extrapolation

(1)  $T_1(q^2) + T_2(q^2)$


(2)  $[T_1+T_2](q^2)$

gives consistent results



# Results for **DI**: $q^2$ -dependence

$T_1, T_2$  (for u/d)

$T_1(q^2)$  and  $T_2(q^2)$  for up or down quark: Disconnected Insertion



$-q^2 a^2$



Slope  $\leftrightarrow$  Signal

$$\sum_{t_1} [\text{Ratio}] = \text{const.} + t_2 \times \underline{[T_{1,2}]}$$

$T_1$

$T_2$


$T_1 + T_2$

$m\pi = 0.48\text{GeV}$



# Results for **Glue**: $q^2$ -dependence

$T_1, T_2$  (for glue)


$T_1(q^2)$  and  $T_2(q^2)$  for glue



$-q^2 a^2$



$a_1 T_1(q^2) + a_2 T_2(q^2)$  for glue



Slope  $\leftrightarrow$  Signal

$T_1$


$T_2$

$m\pi = 0.48\text{GeV}$


$T_1 + T_2$

# Chiral Extrapolation

$T_1, T_2$  (DI) (for u/d)



$T_1, T_2$  (for glue)



Simple Linear-extrapolation is performed

# Renormalization

- Quark-gluon mixing

$$\begin{pmatrix} \langle x \rangle_q^{\overline{MS}}(\mu) \\ \langle x \rangle_G^{\overline{MS}}(\mu) \end{pmatrix} = \begin{pmatrix} Z_{qq}(a\mu, g_0) & Z_{qG}(a\mu, g_0) \\ Z_{Gq}(a\mu, g_0) & Z_{GG}(a\mu, g_0) \end{pmatrix} \begin{pmatrix} \langle x \rangle_q^{lat} \\ \langle x \rangle_G^{lat} \end{pmatrix}$$

Check on Momentum sum rules for lat results

$$\langle x \rangle_q^{lat} + \langle x \rangle_G^{lat} = 0.95(7)$$

$$2(J_q^{lat} + J_G^{lat}) = 0.95(9)$$

←  $Z_{qG} = 0$   
(quenched)

$$Z_{qq} = 1 + \frac{g_0^2}{16\pi^2} C_F \left( \frac{8}{3} \log(a^2 \mu^2) + f_{qq} \right), \quad Z_{qg} = -\frac{g_0^2}{16\pi^2} \left( \frac{2}{3} N_f \log(a^2 \mu^2) + f_{qg} \right),$$

$$Z_{gq} = -\frac{g_0^2}{16\pi^2} C_F \left( \frac{8}{3} \log(a^2 \mu^2) + f_{gq} \right), \quad Z_{gg} = 1 + \frac{g_0^2}{16\pi^2} \left( \frac{2}{3} N_f \log(a^2 \mu^2) + f_{gg} \right).$$

Lat PT calc (one-loop)

← M.Glatzmaier, K.-F.Liu, arXiv:1403.7211

$$\begin{array}{ll} f_{qq} = -7.60930 & f_{qG} = 0 \\ f_{Gq} = -2.37600 & f_{GG} = -3.76900 \end{array}$$

$$\frac{1}{\sqrt{X^\dagger X}} = \int_{-\infty}^{\infty} \frac{d\sigma}{\pi} \frac{1}{\sigma^2 + X^\dagger X}$$

(Integral form for glue op.)

# Renormalization

- “Sum-rule improved” version

$$\langle x \rangle_q^{lat,S} + \langle x \rangle_G^{lat,S} = 1 \quad 2(J_q^{lat,S} + J_G^{lat,S}) = 1$$

“normalization-improvement” by imposing sum-rules to account for latt systematics

$$\langle x \rangle_q^{lat,S} = Z_q^L \langle x \rangle_q^{lat} \quad \langle x \rangle_G^{lat,S} = Z_G^L \langle x \rangle_G^{lat} \quad \text{etc.}$$

– We also have to modify matching coeffs

$$\begin{pmatrix} \langle x \rangle_q^{\overline{MS}}(\mu) \\ \langle x \rangle_G^{\overline{MS}}(\mu) \end{pmatrix} = \begin{pmatrix} Z_{qq}(a\mu, g_0) & Z_{qG}(a\mu, g_0) \\ Z_{Gq}(a\mu, g_0) & Z_{GG}(a\mu, g_0) \end{pmatrix} \begin{pmatrix} \langle x \rangle_q^{lat,S} \\ \langle x \rangle_G^{lat,S} \end{pmatrix}$$

“Sum rule constraint”  $Z_{qq} + Z_{Gq} = 1, \quad Z_{Gq} + Z_{GG} = 1$

$$\Rightarrow \tilde{f}_{qq} = \tilde{f}_{Gq} = (f_{qq} + f_{Gq})/2 \quad \tilde{f}_{qG} = \tilde{f}_{GG} = (f_{qG} + f_{GG})/2$$

$$Z = \begin{pmatrix} 0.9641 & 0.0119 \\ 0.0359 & 0.9881 \end{pmatrix}$$

(ad-hoc solution  
w/ ~1% sys err)

(to MSbar  $\mu=2\text{GeV}$ )

# Results

$$\overline{MS}, \mu = 2 \text{ GeV}$$

(Stat. Error Only)

(Normalization in "2J" unit)

	CI(u)	CI(d)	CI(u+d)	DI(u/d)	DI(s)	Glue
$\langle x \rangle$	0.416(40)	0.151(20)	0.567(45)	0.037(7)	0.023(6)	0.334(56)
$T_2(0)$	0.283(112)	-0.217(80)	0.061(22)	-0.002(2)	-0.001(3)	-0.056(52)
$2J$	0.704(118)	-0.070(82)	0.629(51)	0.035(7)	0.022(7)	0.278(76)
$g_A$	0.91(11)	-0.30(12)	0.62(9)	-0.12(1)	-0.12(1)	—
$2L$	-0.21(16)	0.23(15)	0.01(10)	0.16(1)	0.14(1)	—

$$\text{Quark spin}(g_A) = 25(12)\%$$

$$\text{Quark Orbital}(2L) = 47(13)\%$$

$$\text{Glue total} = 28(08)\%$$

$$\leftarrow (CI(u+d) + 2*DI(u/d) + DI(s))$$

**DI part is important**

$$L(u) + L(d) [CI] \sim 0$$



$$J(u) \gg J(d) [CI] \sim 0$$

(observed in other Lat)


From our old results:  
S.-J.Dong et al.,  
PRL75(1995)2096

# Results

<X>





J



# Results

J



# Systematic errors to be explored

- Dynamical quark effect
  - This is quenched calc.
- Uncertainty in (long) chiral extrapolation
  - $m(\pi) = 0.48\text{--}0.65$  GeV in this calc
- Contamination from excited states
  - Sys error could be large (quite common in N on lat)
- Finite volume artifact, discretization artifact
  - $m(\pi) L \gtrsim 4$ ,  $a = 0.11\text{fm}$
- Renormalization
  - Perturbative vs. non-perturbative, etc.

# Comparison

- quark spin

Quenched calc (1995)

$$\Delta\Sigma^{u,d} \text{ (DI)} \simeq \Delta\Sigma^s \text{ (DI)} \simeq -0.12$$

Recent dynamical clac

$$\Delta\Sigma^{u,d} \text{ (DI)} \sim -0.05$$

(Boston, QCDSF, Engelhardt, ETMC,...)

$$\Delta\Sigma^s \text{ (DI)} \sim -0.03$$

$$L = J - \Delta\Sigma/2$$

→ Large orbital mom by large negative DI in quenched

→ Smaller orbital mom by going to full QCD ?

## HOWEVER:

$$g_A^0 = (\Delta u + \Delta d)[CI] + (\Delta u + \Delta d + \Delta s)[DI] \sim 0.25$$

$$g_A^8 = (\Delta u + \Delta d)[CI] + (\Delta u + \Delta d - 2\Delta s)[DI] = 0.579(25)$$

→ Large DI & larger orbital favored ?

Close-Roberts (1993)

SU(3) breaking effect change situation ?

Lattice calc (Lin et al. (2009), Sasaki et al.(2009), Erkol et al. (2010)) suggests small SU(3) breaking

# Summary & Prospects

- The first study of **complete calc** of proton spin
  - Connected (CI), Disconnected (DI) & Glue
  - DI: stochastic method + unbiased subt. w/ HPE
  - Glue: overlap operator to improved S/N
- Quenched calc at heavy quark mass
  - $J(u+d)$ : 70(5)%,  $J(s)$ : 2.2(7)%,  $J(\text{glue})$ : 28(8)%  
where  $L(u+d+s)$ : 47(13)%
- **Future:**
  - Full QCD calc at lighter mass
  - New approach (Ji, Hatta, ...)