

Rho-meson resonance parameters from Lattice QCD

Dehua Guo, Andrei Alexandru

The George Washington University



The 33rd International Symposium on Lattice Field Theory

July 14, 2015

- 1 Symmetries of the elongated box
- 2 Technical details
- 3 Results
- 4 Conclusions

Symmetries of the elongated box

ρ resonance is in $l = 1, J^P = 1^-$ channel for pion-pion scattering. Elongated box method tunes the momentum of the scattering particles on the lattice.





The $SO(3)$ symmetry group reduce to discrete subgroup O_h or D_{4h}

J	O_h	D_{4h}
0	A_1^+	A_1^+
1	F_1^-	$A_2^- \oplus E^-$
2	$E^+ \oplus F_2^+$	$A_1^+ \oplus B_1^+ \oplus B_2^+ \oplus E^+$
3	$A_2^- \oplus F_1^- \oplus F_2^-$	$A_2^- \oplus B_1^- \oplus B_2^- \oplus 2E^-$
4	$A_1^+ \oplus E^+ \oplus F_1^+ \oplus F_2^+$	$2A_1^+ \oplus A_2^+ \oplus B_1^+ \oplus B_2^+ \oplus 2E^+$

For the p-wave ($l = 1$) scattering channel, we only need to construct the interpolating fields in F_1^- in the O_h group, A_2^- representations in D_{4h} group because the energy contribution from angular momenta $l \geq 3$ is negligible.

Lüscher's formula for elongated box [1]

Phase shift for $l = 1$, rest frame ($\mathbf{P} = 0$):

$$A_2^- : \cot \delta_1(k) = \mathcal{W}_{00} + \frac{2}{\sqrt{5}} \mathcal{W}_{20} \quad (1)$$

(2)

$$\mathcal{W}_{lm}(1, q^2, \eta) = \frac{\mathcal{Z}_{lm}(1, q^2, \eta)}{\eta \pi^{\frac{3}{2}} q^{l+1}}; \quad q = \frac{kL}{2\pi}; \quad \eta = \frac{N_{el}}{N} : \text{elongation factor} \quad (3)$$

Zeta function

$$\mathcal{Z}_{lm}(s; q^2, \eta) = \sum_{\vec{n}} \mathcal{Y}_{lm}(\vec{n}) (\mathbf{n}^2 - q^2)^{-s}; \quad \mathbf{n} \in \mathbf{m} \quad (4)$$


Total energy

$$E = 2\sqrt{m^2 + k^2}; \quad k = \sqrt{\left(\frac{E}{2}\right)^2 - m^2} \quad (5)$$

[1] X. Feng, X. Li, and C. Liu, Phys.Rev. D70 (2004) 014505

Lüscher's formula for boost frame

In order to obtain new kinematic region, we boost the resonance along the elongated direction.



$\mathbf{P} \rightarrow$

$$A_2^- : \cot \delta_1(k) = \mathcal{W}_{00} + \frac{2}{\sqrt{5}} \mathcal{W}_{20} \quad (6)$$

(7)

$$\mathcal{W}_{lm}(1, q^2, \eta) = \frac{\mathcal{Z}_{lm}^{\mathbf{P}}(1, q^2, \eta)}{\gamma \eta \pi^{\frac{3}{2}} q^{l+1}}; \quad \eta = \frac{N_{el}}{N} : \text{elongation factor}; \quad \gamma : \text{boost factor}; \quad (8)$$

$$\mathcal{Z}_{lm}^{\hat{\mathbf{P}}}(s; q^2, \eta) = \sum_{\mathbf{n}} \mathcal{Y}_{lm}(\hat{\mathbf{n}}) (\hat{\mathbf{n}}^2 - q^2)^{-s}; \quad \mathbf{n} \in \frac{1}{\gamma} (\mathbf{m} + \frac{\hat{\mathbf{P}}}{2}); \quad (9)$$



Interpolating field construction for ρ resonance

Four $q\bar{q}$ operators and two scattering operators $\pi\pi$ in A_2^- sector.

$$\rho^J(t_f) = \bar{u}(t_f)\Gamma_{t_f}A_{t_f}(\mathbf{p})d(t_f); \quad \rho^{J\dagger}(t_i) = \bar{d}(t_i)\Gamma_{t_i}^\dagger A_{t_i}^\dagger(\mathbf{p})u(t_i) \quad (10)$$

N	Γ_{t_f}	A_{t_f}	$\Gamma_{t_i}^\dagger$	$A_{t_i}^\dagger$
1	γ_i	$e^{i\mathbf{p}}$	$-\gamma_i$	$e^{-i\mathbf{p}}$
2	$\gamma_4\gamma_i$	$e^{i\mathbf{p}}$	$\gamma_4\gamma_i$	$e^{-i\mathbf{p}}$
3	γ_i	$\nabla_j e^{i\mathbf{p}} \nabla_j$	γ_i	$\nabla_j^\dagger e^{-i\mathbf{p}} \nabla_j^\dagger$
4	$\frac{1}{2}$	$\{e^{i\mathbf{p}}, \nabla_i\}$	$-\frac{1}{2}$	$\{e^{-i\mathbf{p}}, \nabla_i\}$

$$(\pi\pi)_{\mathbf{P}, \Lambda, \mu} = \sum_{\mathbf{p}_1^*, \mathbf{p}_2^*} C(\mathbf{P}, \Lambda, \mu; \mathbf{p}_1; \mathbf{p}_2) \pi(\mathbf{p}_1) \pi(\mathbf{p}_2), \quad (11)$$



$$\pi\pi_{100}(\mathbf{p}_1, \mathbf{p}_2, t) = \frac{1}{\sqrt{2}} [\pi^+(\mathbf{p}_1)\pi^-(\mathbf{p}_2) - \pi^+(\mathbf{p}_2)\pi^-(\mathbf{p}_1)]; \quad \mathbf{p}_1 = (1, 0, 0) \quad \mathbf{p}_2 = (-1, 0, 0)$$

$$\pi\pi_{110} = \frac{1}{2} (\pi\pi(110) + \pi\pi(101) + \pi\pi(1-10) + \pi\pi(10-1))$$

6 × 6 correlation matrix

$$C = \begin{pmatrix} C_{\rho^J \leftarrow \rho^{J'}} & C_{\rho^J \leftarrow \pi \pi_{100}} & C_{\rho^J \leftarrow \pi \pi_{110}} \\ C_{\pi \pi_{100} \leftarrow \rho^{J'}} & C_{\pi \pi_{100} \leftarrow \pi \pi_{100}} & C_{\pi \pi_{100} \leftarrow \pi \pi_{110}} \\ C_{\pi \pi_{110} \leftarrow \rho^{J'}} & C_{\pi \pi_{110} \leftarrow \pi \pi_{100}} & C_{\pi \pi_{110} \leftarrow \pi \pi_{110}} \end{pmatrix}. \quad (12)$$

The correlation functions: $\bar{u}(t_i) \longrightarrow u(t_f)$

$$C_{\rho_i \leftarrow \rho_j} = - \left\langle \begin{array}{c} \Gamma_{t_f}^J(\mathbf{p}, t_f) \\ \text{loop} \\ \Gamma_{t_i}^{J'\dagger}(-\mathbf{p}, t_i) \end{array} \right\rangle = - \langle \text{Tr}[M^{-1}(t_i, t_f) \Gamma_{t_f}^J e^{i\mathbf{p}} M^{-1}(t_f, t_i) \Gamma_{t_i}^{J'\dagger} e^{-i\mathbf{p}}] \rangle. \quad (13)$$

$$C_{\rho_i \leftarrow \pi \pi} = \left\langle \begin{array}{c} \text{triangle} \\ - \\ \text{triangle} \end{array} \right\rangle \stackrel{\mathbf{P} \equiv 0}{=} 2 \left\langle \begin{array}{c} \text{triangle} \end{array} \right\rangle. \quad (14)$$

$$C_{\pi \pi \leftarrow \pi \pi} = - \left\langle \begin{array}{c} \text{square} \\ + \\ \text{square} \\ - \\ \text{X} \\ - \\ \text{X} \\ + \\ \text{figure-eight} \\ - \\ \text{loop} \\ \text{loop} \end{array} \right\rangle \quad (15)$$

$$\stackrel{\mathbf{P} \equiv 0}{=} - \left\langle 2 \begin{array}{c} \text{square} \\ - 2 \text{X} \\ + \text{figure-eight} \\ - \text{loop} \\ \text{loop} \end{array} \right\rangle \quad (16)$$



Laplacian Heaviside smearing [3]



To estimate all-to-all propagators:
The 3-dimensional gauge-covariant Laplacian operator

$$\tilde{\Delta}^{ab}(x, y; U) = \sum_{k=1}^3 \left\{ \tilde{U}_k^{ab}(x) \delta(y, x + \hat{k}) + \tilde{U}_k^{ba}(y)^* \delta(y, x - \hat{k}) - 2\delta(x, y) \delta^{ab} \right\}. \quad (17)$$

$$S_{\Lambda}(t) = \sum_{\lambda(t)}^{\Lambda} |\lambda(t)\rangle \langle \lambda(t)|; \quad \tilde{u}(t) = S(t)u(t) = \sum_{\lambda_t} |\lambda_t\rangle \langle \lambda_t| u(t). \quad (18)$$



[3] C. Morningstar, J. Bulava, J. Foley, K. J. Juge, D. Lenkner, et al., Phys.Rev. D83 (2011)

114505

ρ energy spectrum

We implement the calculation in 3 ensembles ($\eta = 1.0, 1.25, 2.0$) at $m_\pi \approx 310$ MeV and 3 ensembles ($\eta = 1.0, 1.17, 1.33$) at $m_\pi \approx 227$ MeV with nHYP-smearred clover fermions and two mass-degenerated quark flavor.





Figure : The lowest three energy states with their error bars for $\eta = 1.0, m_\pi = 310$ MeV ensemble

We extract energy E by using double exponential $f(t) = we^{-Et} + (1 - w)e^{-E't}$ to do the χ^2 fitting for each eigenvalues.

Energy spectrum



\mathcal{O}_i	1	2	3	4	5	6
	ρ_1	ρ_2	ρ_3	ρ_4	$\pi\pi_{100}$	$\pi\pi_{110}$

(19)

Expectation for energy states





Figure : The lowest 3 energy states prediction from unitary χ PT. When $\eta = 2.0$ the 3rd state is from operator $\pi\pi_{200}$ instead of $\pi\pi_{110}$

Phase shifts and resonance parameters

Figure : Phaseshift data from three ensembles fitted with Breit Wigner form (left) and only fit 5 data points in the resonance region .



$$\cot(\delta_1(E)) = \frac{M_R^2 - E^2}{E\Gamma_r(E)} \text{ where } \Gamma_r(E) \equiv \frac{g_{R12}^2}{6\pi} \frac{p^3}{E^2}. \quad (20)$$

$$\delta_1(E) = \text{arccot} \frac{6\pi(M_R^2 - E^2)E}{g^2 p^3} \quad (21)$$

Centrifugal barrier term [4]

Based on the idea that resonance has finite spatial size, Γ_r is expected to be damped faster than Breit Wigner form above the resonance region. Modify BW with a centrifugal barrier term.

$$\Gamma_r(E) = \frac{g^2}{6\pi} \frac{p^3}{E^2} \frac{1 + (\rho_R R)^2}{1 + (\rho R)^2}. \quad (22)$$




Figure : (left) Current study with LapH smearing vs (right) fitting only the resonance region

[4] F. Von Hippel and C. Quigg, Phys.Rev. D5 (1972) 624-638.

Boost frame data

We add the boost data for the elongated box and fit the phase shift using the modified BW form.




Figure : Left: $m_\rho \approx 310$ MeV. Right: $m_\pi \approx 220$ MeV

m_ρ and $g_{\rho\pi\pi}$ comparison









Figure : m_ρ^2 extrapolation to physical pion mass

Conclusions

- We complete a precision study of ρ resonance with LapH smearing method and obtain the resonance parameters at $m_\pi \approx 310$ MeV and $m_\pi \approx 227$ MeV.
- For precise energy results, Breit Wigner form is not sufficient to fit for the entire region. It needs modification for the BW form.
- The extrapolation of m_ρ to physical pion mass is smaller than $m_\rho^{\text{phy}} = 775$ MeV, we believe that this comes from the absence of strange quark and the $K\bar{K}$ channel which is supported by another unitary χ PT study.

-  X. Feng, X. Li, and C. Liu, *Two particle states in an asymmetric box and the elastic scattering phases*, *Phys.Rev.* **D70** (2004) 014505, [hep-lat/0404001].
-  M. Luscher and U. Wolff, *How to calculate the elastic scattering matrix in two-dimensional quantum field theories by numerical simulation*, *Nucl.Phys.* **B339** (1990) 222–252.
-  C. Morningstar, J. Bulava, J. Foley, K. J. Juge, D. Lenkner, et al., *Improved stochastic estimation of quark propagation with Laplacian Heaviside smearing in lattice QCD*, *Phys.Rev.* **D83** (2011) 114505, [arXiv:1104.3870].
-  F. Von Hippel and C. Quigg, *Centrifugal-barrier effects in resonance partial decay widths, shapes, and production amplitudes*, *Phys.Rev.* **D5** (1972) 624–638.
-  P. Estabrooks and A. D. Martin, *$\pi\pi$ Phase Shift Analysis Below the K anti- K Threshold*, *Nucl.Phys.* **B79** (1974) 301.

Appendix-A: Symmetry on the lattice

The eigenstates $|n\rangle$ are computed in a given irrep of the lattice symmetry group.

$$\psi_n(R^{-1}x) = \psi_n(R^{-1}(x + \mathbf{n}L)); \quad \langle \hat{O}_2(t) \hat{O}_1^\dagger(0) \rangle = \sum_n \langle 0 | \hat{O}_2 | n \rangle \langle n | \hat{O}_1 | 0 \rangle e^{-tE_n} \quad (23)$$

Symmetries: Isospin, flavor, translation, rotation, inversion, etc.

Table : Irreducible representation in $SO(3)$, O and D_4

	$SO(3)$	cubic box(O_h)	elongated box(D_{4h})
irep label	$Y_{lm}; l = 0, 1, \dots, \infty$	A_1, A_2, E, F_1, F_2	A_1, A_2, E, B_1, B_2
dim	$1, 3, \dots, 2l + 1, \dots, \infty$	$1, 1, 2, 3, 3$	$1, 1, 2, 2, 2$

Table : Angular momentum mixing among the irreducible representations of the lattice group

O_h		D_{4h}	
irreducible representation	l	irreducible representation	l
A_1	0, 4, 6, ...	A_1	0, 2, 3, ...
A_2	3, 6, ...	A_2	1, 3, 4, ...
F_1	1, 3, 4, 5, 6, ...	B_1	2, 3, 4, ...
F_2	2, 3, 4, 5, 6, ...	B_2	2, 3, 4, ...
E	2, 4, 5, 6, ...	E	1, 2, 3, 4, ...

Appendix-A: Symmetry on the lattice

The $SO(3)$ symmetry group reduce to discrete subgroup O_h or D_{4h}

Table : Resolution of $2J + 1$ spherical harmonics into the irreducible representations of O_h and D_{4h}

J	O_h	D_{4h}
0	A_1^+	A_1^+
1	F_1^-	$A_2^- \oplus E^-$
2	$E^+ \oplus F_2^+$	$A_1^+ \oplus B_1^+ \oplus B_2^+ \oplus E^+$
3	$A_2^- \oplus F_1^- \oplus F_2^-$	$A_2^- \oplus B_1^- \oplus B_2^- \oplus 2E^-$
4	$A_1^+ \oplus E^+ \oplus F_1^+ \oplus F_2^+$	$2A_1^+ \oplus A_2^+ \oplus B_1^+ \oplus B_2^+ \oplus 2E^+$

Assume that the energy contribution from angular momenta $l \geq 3$ is negligible. For example, if we study the p-wave ($l = 1$) scattering channel, we should construct the interpolating field in F_1^- in the O_h group, A_2^- and E^- representations in D_{4h} group.

Variational method [2]


Variational method is used to extract energy of the excited states.
Construct correlation matrix in the interpolator basis

$$C(t)_{ij} = \langle \mathcal{O}_i(t) \mathcal{O}_j^\dagger(0) \rangle; i, j = 1, 2, \dots, \text{number of operators} \quad (24)$$

The eigenvalues of the correlation matrix are

$$\lambda^{(n)}(t, t_0) \propto e^{-E_n t} (1 + \mathcal{O}(e^{-\Delta E_n t})), n = 1, 2, \dots, \text{number of operators} \quad (25)$$

where $\Delta E_n = E_{\text{Number of operators} + 1} - E_n$.



Larger energy gap makes the high lying energy decay faster and effective mass plateau appear in an earlier time slice.

Appendix-B: LapH smearing

Benefit from LapH smearing:

- Keep low frequency mode up to Λ cutoff to compute the all to all propagators, $u(x) \longrightarrow u(y)$. The number of propagators $M^{-1}(t_f, t_i)$ need to be computed reduce from 6.34×10^{13} in position space to 3.7×10^8 in momentum space for the $24^3 48$ ensemble.
- The effective mass reach a plateau in an earlier time slice.




Figure : pion effective mass with (red) and without LapH smearing (blue)

Appendix-C: Fitting phase-shift




Figure : χ^2 fitting for the phase shift data to Breit Wigner form

$$\chi^2 = \Delta^T COV^{-1} \Delta \quad (26)$$

where

$$\Delta_i = \sqrt{s_i^{\text{curve}}} - \sqrt{s_i^{\text{data}}} \quad (27)$$

Appendix-D: Experiment data [5]




Figure : $\pi\pi$ phase shift below $K\bar{K}$ threshold in experiment

[5] Estabrooks, P. and Martin, Alan D. Nucl.Phys. B79 (1974) 301

$$\Gamma_{BW}(E) = \frac{g^2}{6\pi} \frac{p^3}{E^2}$$

$$\Gamma_{CF}(E) = \frac{g^2}{6\pi} \frac{p^3}{E^2} \frac{1 + (p_R R)^2}{1 + (pR)^2}$$

$$\Gamma_{GA}(E) = \frac{g^2}{6\pi} \frac{p^3}{E^2} \frac{e^{-p^2/6\beta^2}}{e^{-p_R^2/6\beta^2}}$$