



Contribution ID: 256

Type: **Talk**

Charged particles in QED with C^* boundary conditions II

Thursday, 16 July 2015 11:40 (20 minutes)

In order to calculate QED corrections to hadronic quantities by means of lattice simulations, a coherent description of electrically-charged states in finite volume is needed. In the usual periodic setup, Gauss' law and large gauge transformations forbid the propagation of electrically-charged states. A possible solution to this problem, which does not violate the axioms of local quantum field theory, has been proposed by Wiese and Polley, and is based on the use of C^* boundary conditions.

We discuss the properties and symmetries of QED in isolation and QED coupled to QCD, with C^* boundary conditions. We show that a certain class of electrically-charged states can be constructed in this setup in a fully consistent fashion, without relying on gauge fixing. This class of states covers most of the interesting phenomenological applications. We also calculate finite-volume corrections to the mass of stable charged particles and show that these are much smaller than in non-local formulations of QED.

This is the second of two consecutive talks on the subject.

Primary authors: Dr PATELLA, Agostino (CERN and Plymouth University); Dr RAMOS, Alberto (CERN); Prof. LUCINI, Biagio (Swansea University); Dr TANTALO, Nazario (Rome University and INFN "Tor Vergata")

Presenter: Dr TANTALO, Nazario (Rome University and INFN "Tor Vergata")

Session Classification: Theoretical Developments

Track Classification: Theoretical Developments