Real-time calculation of response function with TDHF+BCS

Thursday, 3 April 2008 19:00 (10 minutes)

We present a theoretical study of nuclear response using a method that is applicable to systems from light to heavy nuclei systematically; The real-time calculation of the time-dependent Hartree-Fock+BCS(TDHF+BCS) in the 3-dimensional mesh representation. The method is a simple extension of the well-known TDHF method but is able to describe various responses of deformed nuclei with pairing effects. We show results of our first application to isoscalar monopole and quadrupole modes of excitation for even-even nuclei restricted to N=Z, from Carbon 12 to Zinc 60. Coupling between monopole and quadrupole giant resonances in deformed nuclei is clearly seen in our systematic analysis of the monopole resonance.

Primary author: Mr EBATA, Shuichiro (Institute for Physics, Univ. of Tsukuba)

Co-authors: Prof. YABANA, Kazuhiro (Institute for Physics, Univ. of Tsukuba); Dr NAKATSUKASA, Takashi (Theoretical Nuclear Physics Laboratory, RIKEN Nishina Center); Dr INAKURA, Tsunenori (Institute for Physics, Univ. of Tsukuba); Dr HASHIMOTO, Yukio (Institute for Physics, Univ. of Tsukuba)

Presenter: Mr EBATA, Shuichiro (Institute for Physics, Univ. of Tsukuba)

Session Classification: Poster

Track Classification: Collectivities and shell effects in neutron/proton-rich nuclei