

Observation/Confirmation of hindered E2 strengths in ^{16}C , ^{18}C

Saturday, 5 April 2008 11:00 (20 minutes)

The reduced E2 transition probability from the first excited $2^+ (2^+_{1st})$ state to the ground state — $B(E2)$ — of an even-even nucleus is an important observable that reflects proton collectivity. Recently, an anomalously small $B(E2)$ was reported for the neutron-rich ^{16}C nucleus [1]. The result points to a suppressed proton contribution to the transition strength. This finding raises an intriguing question as to whether or not the neutron contribution is similarly small for the relevant quadrupole excitation. Results from two subsequent experiments [2,3] suggest that the neutron contribution is “normal”. When combined with the $B(E2)$ value, the results indicate a neutron-dominant quadrupole excitation in ^{16}C .

To shed light on the exotic phenomenon exhibited by ^{16}C and to explore the structural evolution of the carbon isotopes towards the neutron dripline, we have carried out an experiment to determine the $B(E2)$ value for the neighboring ^{18}C nucleus.

To determine the $B(E2)$ value for ^{18}C , we have performed an experiment to measure the mean lifetime of the 2^+_{1st} in ^{18}C using an upgraded recoil shadow method (RSM) [1]. The experiment was performed at the RIKEN Nishina Center using the RIPS beamline [4]. Besides ^{18}C , we have also remeasured the mean lifetime of the 2^+_{1st} in ^{16}C . The $B(E2)$ values for ^{16}C , ^{18}C thus determined were about seven and five times smaller than the empirical values, indicating that the anomalously hindered E2 transition observed in ^{16}C persists in ^{18}C . Details of the experiment will be presented and the results will be discussed.

Reference

- [1] N. Imai et al., Phys. Rev. Lett. 92 (2004) 062501.
- [2] Z. Elekes et al., Phys. Lett. B586 (2004) 34.
- [3] H. J. Ong et al., Phys. Rev. C 73 (2006) 024610.
- [4] T. Kubo et al., Nucl. Instrum. Methods B73 (1992) 309.

Primary author: Dr ONG, Hooi Jin (RCNP, Osaka University)

Co-authors: Mr SUZUKI, Daisuke (Univ. of Tokyo); Dr BABA, Hidetada (RIKEN Nishina Center); Dr IWASAKI, Hironori (Univ. of Tokyo); Prof. SAKURAI, Hiroyoshi (RIKEN Nishina Center); Prof. KURITA, Kazuo (Rikkyo Univ.); Dr SUZUKI, Masaru (Univ. of Tokyo); Prof. ISHIHARA, Masayasu (RIKEN Nishina Center); Dr IMAI, Nobuaki (KEK); Dr AOI, Nori (RIKEN Nishina Center); Dr TAKEUCHI, Satoshi (RIKEN Nishina Center); Dr BISHOP, Shawn (RIKEN Nishina Center); Dr OTA, Shunsuke (CNS, Univ. of Tokyo); Prof. NAKAMURA, Takashi (Titech); Dr ONISHI, Takeo (Univ. of Tokyo); Mr NAKAO, Taro (Univ. of Tokyo); Prof. MOTOBAYASHI, Tohru (RIKEN Nishina Center); Mr OKUMURA, Toshifumi (Titech); Prof. KUBO, Toshiyuki (RIKEN Nishina Center); Dr TOGANO, Yasuhiro (Rikkyo Univ.); Dr YANAGISAWA, Yoshiyuki (RIKEN Nishina Center); Dr KONDO, Yosuke (RIKEN Nishina Center); Dr ICHIKAWA, Yuichi (Univ. of Tokyo)

Presenter: Dr ONG, Hooi Jin (RCNP, Osaka University)

Session Classification: Exotic Deformation

Track Classification: Collectivities and shell effects in neutron/proton-rich nuclei