

Extraction of the $\pi^+\pi^-$ subsystem in Diffractively Produced $\pi^-\pi^+\pi^-$ at COMPASS

Friday, 29 July 2016 15:50 (30 minutes)

The COMPASS experiment at CERN has collected a very large data set of 50 million diffractively produced $\pi^-\pi^+\pi^-$ events using a 190 GeV/ c negatively charged hadron beam. The partial-wave analysis (PWA) of these high-precision data reveals previously unseen details. The PWA, which is currently limited by systematic uncertainties, is based on the isobar model, in which multi-particle decays are described as a chain of subsequent two-body decays. In this approach, fixed amplitudes for the intermediate two-pion resonances have to be assumed. These isobar amplitudes are usually parametrized e.g. by Breit-Wigner amplitudes, which enter as prior knowledge, thus increasing systematic uncertainties. We present a novel method, which allows to extract isobar amplitudes directly from the data in a more model-independent way. The focus lies especially on the scalar $\pi^+\pi^-$ subsystem, where in a previous analysis a signal in the $\pi^-\pi^+\pi^-$ sample for a new axial-vector state $a_1(1420)$ in the $f_0(980)\pi$ decay mode was found.

Presenter: KRINNER, Fabian (Technische Universität München Physik Department E18)

Session Classification: Mesons