


# *PT*-symmetric quantum theory, nonlinear eigenvalue problems, and the Painlevé transcendents



The real reason dinosaurs became extinct

**Carl M. Bender**  
*Washington University*

***RIMS-iTHEMS***  
*International Workshop  
on Resurgence Theory  
Kobe, September 2017*

The idea of ***PT***-symmetric quantum theory:


Replace the mathematical condition of Hermiticity by the *weaker* and *physical* condition of ***PT*** symmetry, where

***P*** = *parity*, ***T*** = *time reversal*

(*Physical* because ***P*** and ***T*** are elements of the Lorentz group.)

# A class of $PT$ -symmetric Hamiltonians:

$$H = p^2 + x^2(ix)^\varepsilon \quad (\varepsilon \text{ real})$$



**Look! The energies are real, positive, and discrete for  $\varepsilon > 0$  (!!)**

$$P: x \rightarrow -x, p \rightarrow -p$$

$$T: x \rightarrow x, p \rightarrow -p, i \rightarrow -i$$

CMB and S. Boettcher  
*Physical Review Letters* **80**, 5243 (1998)

# Examples of ***PT***-symmetric Hamiltonians

cubic:  $\varepsilon = 1$

$$H = p^2 + ix^3$$

These Hamiltonians  
have ***PT*** symmetry!


quartic:  $\varepsilon = 2$

$$H = p^2 - x^4$$

← *An upside-down potential with  
real positive eigenvalues!*



THE SPECTRUM OF  $H = p^2 + x^2(ix)^\epsilon$   
IS DISCRETE, REAL, AND  
POSITIVE, AND PARITY  
SYMMETRY IS BROKEN ( $\epsilon > 0$ )



Proof of real eigenvalues:


“ODE/IM Correspondence”  
P. Dorey, C. Dunning, and R. Tateo,  
*J. Phys. A* **40**, R205 (2007)

# ***PT*** symmetry *controls instabilities*


Physical systems that you might *think* are unstable become *stable* in the complex domain...



THE SPECTRUM OF  $H = p^2 + x^2(ix)^\epsilon$   
 IS DISCRETE, REAL, AND  
 POSITIVE, AND PARITY  
 SYMMETRY IS BROKEN IF  $\epsilon > 0$



HEY! WHAT  
 ABOUT  $\epsilon = 2$  ??!



*Upside-down potential with  
 real positive eigenvalues?!*

$$V(x) = -x^4$$

Z. Ahmed, CMB, and M. V. Berry,  
*J. Phys. A: Math. Gen.* **38**, L627 (2005)  
 [arXiv: quant-ph/0508117]

CMB, D. C. Brody, J.-H. Chen, H. F. Jones,  
 K. A. Milton, and M. C. Ogilvie,  
*Phys. Rev. D* **74**, 025016 (2006)  
 [arXiv: hep-th/0605066]

Donald Trump believes in *PT*...

**PUTIN**  
**TRUMP**  
MAKING RUSSIA GREAT AGAIN!

*P* and *T* fit together nicely...









VS KOMT GEISOLEERD UIT G20...



LECTRR

## Stability of the Higgs vacuum:

"***PT***-symmetric interpretation of unstable effective potentials"

CMB, D. W. Hook, N. E. Mavromatos, and S. Sarkar

*Journal of Physics A* **49**, 45LT01 (2016) [arXiv: 1506.01970]

## Stability of the double-scaling limit in QM and QFT:

"***PT***-symmetric Interpretation of double-scaling"

CMB, M. Moshe, and S. Sarkar

*Journal of Physics A* **46**, 102002 (2013) [arXiv: 1206.4943]

"Double-scaling limit of the  $O(N)$ -symmetric anharmonic oscillator"

CMB and S. Sarkar

*Journal of Physics A* **46**, 442001 (2013) [arXiv: 1307.4348]



# *PT* in Asia





# *PT* in Asia



And now  
for something  
completely different...



*Instabilities associated  
with *nonlinear*  
eigenvalue problems...*

**CMB, A. Fring, Q. Wang, and J. Komijani**

# Linear eigenvalue problems...

$$-\psi''(x) + V(x)\psi(x) = E\psi(x)$$

$$\psi(\pm\infty) = 0$$



# For *linear* problems *WKB* gives a good approximation for **large** eigenvalues

$$\int_{x_1}^{x_2} dx \sqrt{E_n - V(x)} \sim (n + 1/2)\pi \quad (n \rightarrow \infty)$$

*n*th energy level grows like a *constant* times a power of *n*

## Example 1: harmonic oscillator

$$V(x) = x^2$$

$$E_n \sim n \quad (n \rightarrow \infty)$$

## Example 2: anharmonic oscillator

$$V(x) = x^4$$

$$E_n \sim Bn^{4/3} \quad (n \rightarrow \infty)$$

$$B = \left[ \frac{3\Gamma(3/4)\sqrt{\pi}}{\Gamma(1/4)} \right]^{4/3}$$

**WKB works for *PT*-symmetric Hamiltonians as well:**

$$H = p^2 + x^2(ix)^\varepsilon \quad (\varepsilon \text{ real})$$

$$E_n \sim \left[ \frac{\Gamma\left(\frac{3}{2} + \frac{1}{\varepsilon+2}\right) \sqrt{\pi} n}{\sin\left(\frac{\pi}{\varepsilon+2}\right) \Gamma\left(1 + \frac{1}{\varepsilon+2}\right)} \right]^{\frac{2\varepsilon+4}{\varepsilon+4}} \quad (n \rightarrow \infty)$$

# Asymptotics beyond all orders

Leading asymptotic behavior of solutions to

$$-\psi''(x) + V(x)\psi(x) = E\psi(x)$$

for large positive  $x$ :

$$\psi(x) \sim C[V(x) - E]^{-1/4} \exp \left[ \int^x ds \sqrt{V(s) - E} \right] \quad (x \rightarrow \infty)$$


**NOTE:** There is only **ONE** arbitrary constant.

Second arbitrary constant is invisible with Poincaré asymptotics because it is contained in the *subdominant* solution:

$$\psi(x) \sim D[V(x) - E]^{-1/4} \exp \left[ - \int^x ds \sqrt{V(s) - E} \right] \quad (x \rightarrow \infty)$$

*Physical* solution is **Unstable** under small changes in  $E$ .

# Eigenfunctions: 3 characteristic properties



- (1) **Oscillatory** in *classically allowed* region ( $n$ th eigenfunction has  $n$  nodes)
- (2) **Monotone decay** in *classically forbidden* region
- (3) **Transition** at the boundary (*turning point*)




# Toy nonlinear eigenvalue problem

$$y'(x) = \cos[\pi xy(x)], \quad y(0) = a$$


## Some references:

- [1] C. M. Bender and S. A. Orszag, *Advanced Mathematical Methods for Scientists and Engineers* (McGraw Hill, New York, 1978), chap. 4.
- [2] C. M. Bender, D. W. Hook, P. N. Meisinger, and Q. Wang, *Phys. Rev. Lett.* **104**, 061601 (2010).
- [3] C. M. Bender, D. W. Hook, P. N. Meisinger, and Q. Wang, *Ann. Phys.* **325**, 2332-2362 (2010).
- [4] J. Gair, N. Yunes, and C. M. Bender, *J. Math. Phys.* **53**, 032503 (2012).



**Solutions for 50 initial conditions**

**Note: (1) oscillation (2) monotone decay (3) transition**




# Asymptotic behavior for large $x$

**Solution behaves like:**  $y(x) \sim \frac{m + 1/2}{x}$

$m = 0, 1, 2, 3, \dots$  is an integer

There's a ***big*** problem here...



Where are the **odd- $m$**  solutions?!?



**Furthermore, no arbitrary constant appears  
in the asymptotic behavior!!**



# Where is the arbitrary constant?!?



Is it in higher order?

# Higher-order asymptotic behavior for large $x$ still contains no arbitrary constant!

$$y(x) \sim \frac{m + 1/2}{x} + \sum_{k=1}^{\infty} \frac{c_k}{x^{2k+1}} \quad (x \rightarrow \infty)$$

$$c_1 = \frac{(-1)^m}{\pi}(m + 1/2),$$

$$c_2 = \frac{3}{\pi^2}(m + 1/2),$$

$$c_3 = (-1)^m \left[ \frac{(m + 1/2)^3}{6\pi} + \frac{15(m + 1/2)}{\pi^3} \right],$$

$$c_4 = \frac{8(m + 1/2)^3}{3\pi^2} + \frac{105(m + 1/2)}{\pi^4},$$

$$c_5 = (-1)^m \left[ \frac{3(m + 1/2)^5}{40\pi} + \frac{36(m + 1/2)^3}{\pi^3} + \frac{945(m + 1/2)}{\pi^5} \right],$$


$$c_6 = \frac{38(m + 1/2)^5}{15\pi^2} + \frac{498(m + 1/2)^3}{\pi^4} + \frac{10395(m + 1/2)}{\pi^6}.$$

# Asymptotics beyond all orders

**Difference of two solutions in one bundle:**  $Y(x) \equiv y_1(x) - y_2(x)$


$$\begin{aligned} Y'(x) &= \cos[\pi x y_1(x)] - \cos[\pi x y_2(x)] \\ &= -2 \sin \left[ \frac{1}{2} \pi x y_1(x) + \frac{1}{2} \pi x y_2(x) \right] \sin \left[ \frac{1}{2} \pi x y_1(x) - \frac{1}{2} \pi x y_2(x) \right] \\ &\sim -2 \sin \left[ \pi \left( m + \frac{1}{2} \right) \right] \sin \left[ \frac{1}{2} \pi x Y(x) \right] \quad (x \rightarrow \infty) \\ &\sim -(-1)^m \pi x Y(x) \quad (x \rightarrow \infty). \end{aligned}$$

$$Y(x) \sim K \exp \left[ -(-1)^m \pi x^2 \right] \quad (x \rightarrow \infty)$$



**Aha!  $K$  is the invisible arbitrary constant!**  
**Odd- $m$  solutions are *unstable*;**  
**even- $m$  solutions are *stable*.**





$m = 9$   
 $m = 7$   
 $m = 5$   
 $m = 3$   
 $m = 1$

$$y(0) = a \in \{1.6026, 2.3884, 2.9767, 3.4675, 3.8975, 4.2847, \dots\}$$

Eigenvalues correspond to **odd- $m$**  initial values.  
Eigenfunctions are (*unstable*) **separatrices**, which begin at eigenvalues.

# We calculated up to $m=500,001$

Let  $m = 2n - 1$

For large  $n$  the  $n$ th eigenvalue grows like the *square root* of  $n$  times a constant  $A$ , and we used Richardson extrapolation to show that

$A = 1.7817974363\dots$

and then we guessed  $A$ .



# Result:



$$a_n \sim A\sqrt{n} \quad (n \rightarrow \infty)$$

$$A = 2^{5/6}$$

**This is a rather nontrivial problem...**

# Analytic calculation of the constant $A$

Construct moments of  $z(t)$ :

$$A_{n,k}(t) \equiv \int_0^t ds \cos[n\lambda s z(s)] \frac{s^{k+1}}{[z(s)]^k}$$

Moments are associated with a semi-infinite **linear** one-dimensional random walk in which random walkers become static as they reach  $n=1$

$$2\alpha_{1,k} + \alpha_{2,k-1} = 0, \quad 2\alpha_{n,k} + \alpha_{n-1,k-1} + \alpha_{n+1,k-1} = 0 \quad (n \geq 3).$$

$$2\alpha_{2,k} + \alpha_{3,k-1} = 0,$$

Solve the random walk problem exactly and get  $A = 2^{5/6}$



CMB, A. Fring, and J. Komijani  
*J. Phys. A: Math. Theor.* **47**, 235204 (2014)  
[arXiv: math-ph/1401.6161]



# Possible connection with the *power series constant* $P$ ???

(Remember the numerical constant  $A = 1.7818$ )

W. K. Hayman, *Research Problems in Function theory*  
[Athlone Press (University of London), London, 1967]

J. Clunie and P. Erdős, *Proc. Roy. Irish Acad.* **65**, 113 (1967).  
J. D. Buckholtz, *Michigan Math. J.* **15**, 481 (1968).

$$1 \leq P \leq 2$$

$$\sqrt{2} \leq P \leq 2$$

$$1.7 \leq P \leq 12^{1/4}$$

$$1.7818 \leq P \leq 1.82$$

# *Three nontrivial second-order nonlinear eigenvalue problems*



**separatrix**

# Painlevé equations



**Paul Painlevé  
(1863-1933)**

**Six Painlevé equations known as Painlevé I – VI**

**Only spontaneous singularities are poles**

## Painlevé I

$$\frac{d^2y}{dt^2} = 6y^2 + t$$

## Painlevé II

$$\frac{d^2y}{dt^2} = 2y^3 + ty + \alpha$$

## Painlevé III

$$ty \frac{d^2y}{dt^2} = t \left( \frac{dy}{dt} \right)^2 - y \frac{dy}{dt} + \delta t + \beta y + \alpha y^3 + \gamma ty^4$$

## Painlevé IV

$$y \frac{d^2y}{dt^2} = \frac{1}{2} \left( \frac{dy}{dt} \right)^2 + \beta + 2(t^2 - \alpha)y^2 + 4ty^3 + \frac{3}{2}y^4$$

## Painlevé V

$$\begin{aligned} \frac{d^2y}{dt^2} &= \left( \frac{1}{2y} + \frac{1}{y-1} \right) \left( \frac{dy}{dt} \right)^2 - \frac{1}{t} \frac{dy}{dt} \\ &+ \frac{(y-1)^2}{t^2} \left( \alpha y + \frac{\beta}{y} \right) + \gamma \frac{y}{t} + \delta \frac{y(y+1)}{y-1} \end{aligned}$$

## Painlevé VI

$$\begin{aligned} \frac{d^2y}{dt^2} &= \frac{1}{2} \left( \frac{1}{y} + \frac{1}{y-1} + \frac{1}{y-t} \right) \left( \frac{dy}{dt} \right)^2 - \left( \frac{1}{t} + \frac{1}{t-1} + \frac{1}{y-t} \right) \frac{dy}{dt} \\ &+ \frac{y(y-1)(y-t)}{t^2(t-1)^2} \left( \alpha + \beta \frac{t}{y^2} + \gamma \frac{t-1}{(y-1)^2} + \delta \frac{t(t-1)}{(y-t)^2} \right) \end{aligned}$$



# (1) First Painlevé transcendent

$$y''(t) = 6[y(t)]^2 + t, \quad y(0) = b, \quad y'(0) = c$$

Solution  $y(x)$  must *choose* between two possible asymptotic behaviors as  $x$  gets large and negative:

$$+\sqrt{-t/6} \text{ or } -\sqrt{-t/6}.$$

# Example of a *difficult* choice ...



# Two possible asymptotic behaviors

Lower square-root branch is *stable*:


$$y(x) \sim -\sqrt{-x} + c(-x)^{-1/8} \cos \left[ \frac{4}{5} \sqrt{2} (-x)^{5/4} + d \right] \quad (x \rightarrow -\infty)$$

Upper square-root branch is *unstable*:


$$y(x) \sim \sqrt{-x} + c_{\pm} (-x)^{-1/8} \exp \left[ \pm \frac{4}{5} \sqrt{2} (-x)^{5/4} \right] \quad (x \rightarrow -\infty)$$

# Two possible kinds of solutions (NOT eigenfunctions):


**Unstable branch**




**Stable branch**



**Unstable branch**





**Stable branch**



# First four separatrix (eigenfunction) solutions:



Unstable

Stable



Unstable



Stable



*Initial slope is the eigenvalue, initial value  $y(0) = 0$*







# Tenth and eleventh separatrix (eigenfunction) solutions:



*Initial slope* is the eigenvalue, initial value  $y(0) = 0$

# First four separatrix solutions with 0 initial slope:



# Numerical calculation of eigenvalues

(*nonlinear* semiclassical large- $n$  limit)

$$y'(0) = b_n \quad y(0) = 0$$

$$b_n \sim B_I n^{3/5} \quad B_I = 2.09214674$$

$$y(0) = c_n \quad y'(0) = 0$$

$$c_n \sim C_I n^{2/5} \quad C_I = -1.0304844$$

$$y_n(0) \sim -1.0304844 \left(n - \frac{1}{2}\right)^{2/5} \left[1 - \frac{0.0096518}{\left(n - \frac{1}{2}\right)^2} + \frac{0.0240}{\left(n - \frac{1}{2}\right)^4}\right]$$

# Analytical asymptotic calculation of eigenvalues

$$B_1 = 2 \left[ \frac{5\sqrt{\pi}\Gamma(5/6)}{2\sqrt{3}\Gamma(1/3)} \right]^{3/5}$$

$$C_1 = - \left[ \frac{5\sqrt{\pi}\Gamma(5/6)}{2\sqrt{3}\Gamma(1/3)} \right]^{2/5}$$

Obtained by using WKB to calculate the large eigenvalues of the  
*cubic  $PT$ -symmetric Hamiltonian*

$$H = \frac{1}{2}p^2 + 2ix^3$$

**Painlevé I corresponds to  $\varepsilon = 1$**

(Do you remember  
the cubic  *$PT$* -symmetric  
Hamiltonian?!)





# Analytical asymptotic calculation of eigenvalues

**Multiply Painlevé I equation by  $y'(t)$ ;**


**Then integrate from  $t = 0$  to  $t = x$ :**

$$H \equiv \frac{1}{2}[y'(x)]^2 - 2[y(x)]^3 = \frac{1}{2}[y'(0)]^2 - 2[y(0)]^3 + I(x),$$

**where  $I(x) = \int_0^x dt ty'(t)$ .**

**Take  $|x|$  large at an angle of  $\pi/4$ ,  $I(x) \rightarrow 0$ , and we get the *PT*-symmetric Hamiltonian for  $\varepsilon = 1$ .**

D. Masoero noted connections between Painlevé I and  $H = p^2 + ix^3$



## (2) Second Painlevé transcendent

$$y''(t) = 2[y(t)]^3 + ty(t), \quad y(0) = b, \quad y'(0) = c$$

Now, both solutions

$$+\sqrt{-t/2} \text{ or } -\sqrt{-t/2}$$



are unstable and  $0$  is stable.

# Two types of solutions (not eigenfunctions):

**Unstable**

**Stable**



**Unstable**







**Unstable**

**Stable**



**Unstable**



# First four separatrix solutions with $y(0)=0$ :





# 20<sup>th</sup> and 21<sup>st</sup> separatrix solutions:







# First four separatrices with vanishing initial slope $y'(0)=0$ :





Unstable



Unstable

# 13<sup>th</sup> and 14<sup>th</sup> separatrices:



# Numerical calculation of eigenvalues

$$y(0) = 0, b_n = y'(0)$$

$$c_n = y(0), y'(0) = 0$$

$$b_n \sim B_{\text{II}} n^{2/3} \quad \text{and} \quad c_n \sim C_{\text{II}} n^{1/3}$$

$$B_{\text{II}} = 1.8624128$$

$$C_{\text{II}} = 1.21581165$$

$$y_n(0) \sim 1.2158117 n^{1/3} \left[ 1 + \frac{0.0052543}{n^2} + \frac{0.077}{n^4} \right]$$

**CMB and J. Komijani**

**J. Physics A: Math. Theor. 48, 475202 (2015)**

# Analytical calculation of eigenvalues

$$B_{II} = \left[ 3\sqrt{2\pi}\Gamma\left(\frac{3}{4}\right) / \Gamma\left(\frac{1}{4}\right) \right]^{2/3}$$

$$C_{II} = \left[ 3\sqrt{\pi}\Gamma\left(\frac{3}{4}\right) / \Gamma\left(\frac{1}{4}\right) \right]^{1/3}$$

Obtained by using WKB to calculate the large eigenvalues of the  
*quartic  $PT$ -symmetric Hamiltonian*

$$H = \frac{1}{2}p^2 - \frac{1}{2}x^4$$

**Painlevé II corresponds to  $\varepsilon = 2$**

(Do you remember the  
quartic upside-down  
 *$PT$* -symmetric  
Hamiltonian?!)



### (3) Fourth Painlevé transcendent

$$y(t)y''(t) = \frac{1}{2}[y'(t)]^2 + 2t^2[y(t)]^2 + 4t[y(t)]^3 + \frac{3}{2}[y(t)]^4$$



with  $y(0) = c$  and  $y'(0) = b$ .



# Two possible kinds of solutions (NOT eigenfunctions):



Unstable branch

Stable branch



Unstable branch



Stable branch



# First four separatrix (eigenfunction) solutions [ $y(0)=1$ ]:



Unstable

Stable





Unstable

Stable



# Tenth and eleventh separatrix (eigenfunction) solutions:





*Slope* is the eigenvalue, initial value  $y(0) = 1$

# First four separatrix (eigenfunction) solutions [ $y'(0)=0$ ]:



Unstable

Stable





Unstable

Stable



# Tenth and eleventh separatrix (eigenfunction) solutions:



**$y(0)$  is the eigenvalue, initial slope is 0**

Large  $n$  behaviour of eigenvalues:  $b_n \sim B_{\text{IV}} n^{3/4}$  and  $c_n \sim C_{\text{IV}} n^{1/2}$ .

Numerical results using Richardson extrapolation:

$$B_{\text{IV}} = 4.256843.$$

$$C_{\text{IV}} = -2.626587.$$

Analytic results using  $\hat{H} = \frac{1}{2}\hat{p}^2 + \frac{1}{8}\hat{x}^6$ .

$$B_{\text{IV}} = 2^{3/2} \left[ \sqrt{\pi} \Gamma\left(\frac{5}{3}\right) / \Gamma\left(\frac{7}{8}\right) \right]^{3/4},$$

$$C_{\text{IV}} = -2 \left[ \sqrt{\pi} \Gamma\left(\frac{5}{3}\right) / \Gamma\left(\frac{7}{8}\right) \right]^{1/2}.$$

Obtained by using WKB to calculate the large eigenvalues of the

***sextic  $PT$ -symmetric Hamiltonian***



**Note:**

**Painlevé I, II, and IV  
correspond to  $\varepsilon = 1, 2,$  and  $4$**



**This analysis extends to huge classes of equations beyond Painlevé. For example:**

**Super Painlevé:**

$$y''(x) = \frac{2M + 2}{(M - 1)^2} [y(x)]^M + x[y(x)]^N$$



$$y'' = \frac{10}{9}y^4 + x$$

$$y(0) = 0$$

$$y'_n(0) \sim 1.1102 n^{\frac{5}{11}} \left(1 + \frac{26.235}{11n}\right)$$

$$y'_n(0) \sim -1.109 n^{\frac{5}{11}} \left(1 - \frac{11}{11n}\right)$$

$$y'(0) = 0$$

$$y_n(0) \sim 1.80547 n^{\frac{2}{11}} \left(1 - \frac{1.998}{11n}\right)$$

$$y_n(0) \sim -1.226 n^{\frac{2}{11}} \left(1 + \frac{3.03}{11n}\right)$$

$$y'' = \frac{10}{9}y^4 + xy$$

$$y(0) = 0$$

$$y'_n(0) \sim 2.1336n^{\frac{5}{9}} \begin{cases} \left(1 - \frac{3.55}{9n}\right), & \text{odd } n \\ \left(1 - \frac{2.13}{9n}\right), & \text{even } n \end{cases}$$

$$y'_n(0) \sim -2.1336n^{\frac{5}{9}} \begin{cases} \left(1 - \frac{3.24}{9n}\right), & \text{odd } n \\ \left(1 - \frac{2.08}{9n}\right), & \text{even } n \end{cases}$$

$$y'(0) = 0$$

$$y_n(0) \sim -1.59255n^{\frac{2}{9}} \begin{cases} \left(1 - \frac{1.24}{9n}\right), & \text{odd } n \\ \left(1 - \frac{0.77}{9n}\right), & \text{even } n \end{cases}$$

$$y'' = \frac{10}{9}y^4 + xy^2$$

$$y(0) = 0$$

$$y'_n(0) \sim 2.9996n^{\frac{5}{7}} \left(1 - \frac{0.959}{7n}\right)$$

$$y'_n(0) \sim -2.9996n^{\frac{5}{7}} \left(1 - \frac{3.19}{7n}\right)$$

$$y'(0) = 0$$

$$y_n(0) \sim 1.098102n^{\frac{2}{7}} \left(1 - \frac{2.00207}{7n}\right)$$

$$y_n(0) \sim -1.82502n^{\frac{2}{7}} \left(1 - \frac{0.84}{7n}\right)$$

$$y'' = \frac{14}{25}y^6 + x$$

$$y(0) = 0$$

$$y'_n(0) \sim 2.3219n^{\frac{7}{17}} \left(1 + \frac{2.52}{17n}\right)$$

$$y'_n(0) \sim -2.322n^{\frac{7}{17}} \left(1 - \frac{7.3}{17n}\right)$$

$$y'(0) = 0$$

$$y_n(0) \sim 1.500998n^{\frac{2}{17}} \left(1 - \frac{1.9992}{17n}\right)$$

$$y_n(0) \sim -1.652812n^{\frac{2}{17}} \left(1 - \frac{0.6732}{17n}\right)$$

$$y'' = \frac{14}{25}y^6 + xy$$

$$y(0) = 0$$

$$y'_n(0) \sim 2.3569n^{\frac{7}{15}} \begin{cases} \left(1 + \frac{4.137}{15n}\right), & \text{odd } n \\ \left(1 + \frac{0.4425}{15n}\right), & \text{even } n \end{cases}$$

$$y'_n(0) \sim -2.357n^{\frac{7}{15}} \begin{cases} \left(1 - \frac{4.125}{15n}\right), & \text{odd } n \\ \left(1 - \frac{0.4275}{15n}\right), & \text{even } n \end{cases}$$

$$y'' = \frac{14}{25}y^6 + xy^2$$

$$y(0) = 0$$

$$y'_n(0) \sim 2.5979n^{\frac{7}{13}} \begin{cases} \left(1 - \frac{1.2675}{13n}\right), & \text{odd } n \\ \left(1 - \frac{1.8473}{13n}\right), & \text{even } n \end{cases}$$

$$y'_n(0) \sim -2.598n^{\frac{7}{13}} \begin{cases} \left(1 - \frac{5.148}{13n}\right), & \text{odd } n \\ \left(1 - \frac{5.727}{13n}\right), & \text{even } n \end{cases}$$

$$y'(0) = 0$$

$$y_n(0) \sim 1.73085n^{\frac{2}{13}} \left(1 + \frac{0.195}{13n}\right)$$

$$y_n(0) \sim -1.7065n^{\frac{2}{13}} \begin{cases} \left(1 - \frac{0.913}{13n}\right), & \text{odd } n \\ \left(1 - \frac{1.08}{13n}\right), & \text{even } n \end{cases}$$

$$y'' = \frac{14}{25}y^6 + xy^3$$

$$y(0) = 0$$

$$y'_n(0) \sim 1.7408n^{\frac{7}{11}} \begin{cases} \left(1 + \frac{0.33}{11n}\right), & \text{odd } n \\ \left(1 - \frac{4.58}{11n}\right), & \text{even } n \end{cases}$$

$$y'_n(0) \sim -1.7408n^{\frac{7}{11}} \begin{cases} \left(1 - \frac{2.44}{11n}\right), & \text{odd } n \\ \left(1 - \frac{7.34}{11n}\right), & \text{even } n \end{cases}$$

$$y'(0) = 0$$

$$y_n(0) \sim -1.52224n^{\frac{2}{11}} \begin{cases} \left(1 + \frac{0.305}{11n}\right), & \text{odd } n \\ \left(1 + \frac{1.705}{11n}\right), & \text{even } n \end{cases}$$

$$y'' = \frac{14}{25}y^6 + xy^4$$

$$y(0) = 0, x < 0$$

$$y'_n(0) \sim 3.06787n^{\frac{7}{9}} \begin{cases} \left(1 + \frac{1.5}{9n}\right), & \text{odd } n \\ \left(1 - \frac{1.87}{9n}\right), & \text{even } n \end{cases}$$

$$y'_n(0) \sim -3.06786n^{\frac{7}{9}} \begin{cases} \left(1 - \frac{5.13}{9n}\right), & \text{odd } n \\ \left(1 - \frac{8.51}{9n}\right), & \text{even } n \end{cases}$$

$$y(0) = 0, x > 0$$

$$y'_n(0) \sim 2.9010n^{\frac{7}{9}} \left(1 - \frac{1.67}{9n}\right)$$

$$y'_n(0) \sim -2.9010n^{\frac{7}{9}} \left(1 + \frac{1.65}{9n}\right)$$

$$y'' = \frac{4}{9}y^7 + xy^4$$

$$y(0) = 0, \quad x < 0$$

$$y'_n(0) \sim -1.38115n^{\frac{2}{3}} \left(1 - \frac{0.4635}{3n}\right)$$

$$y(0) = 0, \quad x > 0$$

$$y'_n(0) \sim -1.38114n^{\frac{2}{3}} \left(1 - \frac{0.462}{3n}\right)$$

$$y'' = \frac{4}{9}y^7 + xy^5$$

$$y(0) = 0, \quad x < 0$$

$$y'_n(0) \sim -1.86695n^{\frac{4}{5}} \begin{cases} \left(1 - \frac{0.7375}{5n}\right), & \text{odd } n \\ \left(1 - \frac{3.2575}{5n}\right), & \text{even } n \end{cases}$$

$$y(0) = 0, \quad x > 0$$

$$y'_n(0) \sim -2.29535n^{\frac{4}{5}} \left(1 - \frac{0.225}{5n}\right)$$

# Hyperfine splitting

$$y'' = \frac{1}{a^2}y^4 + xy^2, \quad \text{with} \quad a \equiv \frac{3}{\sqrt{10}}.$$


Let  $y_{nm}(x) = Y_n(x) + \phi_m(x)$ , where  $Y_n(x)$  is a separatrix solution with

$$Y_n(x) \sim a\sqrt{-x}, \quad x \rightarrow -\infty.$$

The new hyperfine solutions initially follow  $Y_n(x)$ .

Then they deviate from  $Y_n(x)$  and oscillate  $m$  times about the curve  $-a\sqrt{-x}$ .

Finally, they level off for large  $x$  as  $y_{nm}(x) \sim \frac{12}{x^3}, \quad x \rightarrow -\infty.$





The initial values of  $\phi$  are the hyperfine eigenvalues.


For example, for the lowest eigenfunction  $Y_0$ ,

$$\phi_m(0) \sim 4.1789 e^{-9.26201m}, \quad m \rightarrow \infty.$$

The hyperfine oscillation separates at the negative values

$$T_m \sim \left( \frac{7}{4\sqrt{2a}} 9.26201m \right)^{\frac{4}{7}}, \quad m \rightarrow \infty.$$

We hope we have opened a window  
to a new area of *nonlinear*  
*semiclassical* asymptotic analysis



Download from  
Dreamstime.com  
This watermarked comp image is for previewing purposes only.



1917393

Miguel angel Salinas salinas | Dreamstime.com

*Thanks for listening!*