

**RIMS-iTHEMS Workshop on Resurgence
6.9.2017@Kobe**

Hofstadter, Toda & Calabi-Yau


**Yasuyuki Hatsuda
(Rikkyo University)**

**RIMS-iTHEMS Workshop on Resurgence
6.9.2017@Kobe**


**Hofstadter, Toda & Calabi-Yau
(2d electrons, integrable systems
& geometry)**

**Yasuyuki Hatsuda
(Rikkyo University)**

In 1976, Douglas Hofstadter
considered an interesting 2d
electron model in a magnetic field



He predicted a novel electron spectrum, which is now known as **Hofstadter's butterfly**



**40 years later, Katsura,
Tachikawa and myself found
that the completely same figure
appears in the context of
Calabi-Yau geometry**

**YH, Katsura & Tachikawa, arXiv:1606.01894
“Hofstadter’s butterfly in quantum geometry”**

**In this talk, I would like to
explain its idea**

Consequence

YH-Katsura-Tachikawa

New J.Phys. 18 (2016) 10, 103023

Moduli of CYs

Density of states

$$\rho(E) = \frac{1}{2\pi} \operatorname{Im} \left[\frac{\partial t(E, q)}{\partial E} \right]$$

Magnetic effect

= “Quantum deformation”

[Recall Dunne’s & Schiappa’s talk]

Contents

1. **Hofstadter** Model
2. **Toda** Lattice
3. Relation to **Calabi-Yau**
4. **Semiclassical Analysis**


1. Hofstadter Model

2d Electron in Magnetic Field

$$H = \frac{1}{2m} (\mathbf{p} + e\mathbf{A})^2 = \frac{1}{2m} (\Pi_x^2 + \Pi_y^2)$$

Vector potential

$$\Pi := \mathbf{p} + e\mathbf{A}$$




2d Electron in Magnetic Field

$$H = \frac{1}{2m} (\mathbf{p} + e\mathbf{A})^2 = \frac{1}{2m} (\Pi_x^2 + \Pi_y^2)$$

Vector potential

$$\mathbf{\Pi} := \mathbf{p} + e\mathbf{A}$$



$$[\Pi_x, \Pi_y] = \frac{\hbar e}{i} (\partial_x A_y - \partial_y A_x) = -i\hbar e B$$


2d Electron in Magnetic Field

- This Hamiltonian is the same as that for the harmonic oscillator

$$E_n = \hbar\omega_c \left(n + \frac{1}{2} \right)$$

$$\omega_c = \frac{eB}{m}$$


Landau level



- The spectrum of the 2d electron is **quantized** by the magnetic effect

An Electron on 2d Lattice

- In this case, the spectrum is obtained by the **tight-binding approximation**



$$E = 2 \cos(k_x a) + 2 \cos(k_y a)$$


- The allowed range of the energy:

$$-4 \leq E \leq 4$$

Single energy band

The Hofstadter Model

- An electron on a 2d lattice with a magnetic flux



$$\text{Free } E = \frac{\hbar^2}{2m} (k_x^2 + k_y^2) \xrightarrow{B} H = \frac{1}{2m} (\Pi_x^2 + \Pi_y^2)$$


Lattice

$$\Pi := p + eA$$

$$E = e^{ik_x a} + e^{-ik_x a} + e^{ik_y a} + e^{-ik_y a}$$

The Hofstadter Model

- An electron on a 2d lattice with a magnetic flux



$$\text{Free } E = \frac{\hbar^2}{2m} (k_x^2 + k_y^2) \xrightarrow{B} H = \frac{1}{2m} (\Pi_x^2 + \Pi_y^2)$$

$$\Pi := p + eA$$

Lattice

$$E = e^{ik_x a} + e^{-ik_x a} + e^{ik_y a} + e^{-ik_y a}$$

$$H = e^{\frac{ia}{\hbar} \Pi_x} + e^{-\frac{ia}{\hbar} \Pi_x} + e^{\frac{ia}{\hbar} \Pi_y} + e^{-\frac{ia}{\hbar} \Pi_y}$$

The Hofstadter Model

- **By fixing the Landau gauge $A = (0, Bx, 0)$, the eigenvalue problem finally leads to**

Harper's equation


Hofstadter '76

$$\psi(n+1) + \psi(n-1) + 2 \cos(2\pi n\alpha + \nu)\psi(n) = E\psi(n)$$

$$\alpha = \frac{a^2 e B}{2\pi \hbar}$$

- **If $\alpha = P/Q$ (rational), the spectrum of this equation gives Q energy bands**

Hofstadter's Butterfly



2. Toda Lattice

The Toda Lattice

- **The Toda lattice is the well-known **quantum integrable system**** **Toda '67**

$$H_1 = \sum_{n=1}^N \left(\frac{p_n^2}{2} + e^{x_n - x_{n+1}} \right), \quad x_{N+1} = x_1$$

- **There are N mutually commuting operators**

$$[H_n, H_m] = 0$$

- **The eigenvalue problem**

$$H_k \Psi(x_1, \dots, x_N) = E_k \Psi(x_1, \dots, x_N)$$

Generalized Toda Lattice

- **There is a one-parameter deformation of the Toda lattice**

Ruijsenaars '90

$$H_1 = \sum_{n=1}^N \left(1 + q^{-1/2} R^2 e^{x_n - x_{n+1}} \right) e^{R p_n}$$

$$q = e^{iR\hbar}$$

Generalized Toda Lattice

- There is a one-parameter deformation of the Toda lattice Ruijsenaars '90

$$H_1 = \sum_{n=1}^N \left(1 + q^{-1/2} R^2 e^{x_n - x_{n+1}} \right) e^{R p_n}$$



$R \rightarrow 0$

$$q = e^{iR\hbar}$$

$$H_1 = N + R \sum_{n=1}^N p_n$$

$$+ R^2 \sum_{n=1}^N \left(\frac{p_n^2}{2} + e^{x_n - x_{n+1}} \right) + \mathcal{O}(R^3)$$

N=2 gToda

- **Let us consider the case of N=2**

$$H = e^{Rp_1} + e^{Rp_2} + R^2 (e^{x_1 - x_2 + Rp_1} + e^{x_2 - x_1 + Rp_2})$$

$$p_1 + p_2 = 0 \quad \textbf{(Center of mass frame)}$$

$$p := Rp_1, \quad x := x_1 - x_2 + Rp_1$$

N=2 gToda

- **Let us consider the case of N=2**

$$H = e^{Rp_1} + e^{Rp_2} + R^2 (e^{x_1 - x_2 + Rp_1} + e^{x_2 - x_1 + Rp_2})$$

$$p_1 + p_2 = 0 \quad \textbf{(Center of mass frame)}$$

$$p := Rp_1, \quad x := x_1 - x_2 + Rp_1$$

$$H = e^p + e^{-p} + R^2 (e^x + e^{-x})$$

$$[x, p] = iR\hbar$$

N=2 gToda

- The eigenvalue equation is thus a **difference equation**

$$\psi(x + iR\hbar) + \psi(x - iR\hbar) + 2R^2 \cosh x \psi(x) = E\psi(x)$$

- This equation is very similar to **Harper's equation**, but their spectra are quite different

Harper → **Continuous (Bands)**

Toda → **Discrete**

3. Relation to Calabi-Yau

So far...

- The **Hofstadter** model (Harper's equation)

$$\psi(n+1) + \psi(n-1) + 2 \cos(2\pi n\alpha + \nu)\psi(n) = E\psi(n)$$

- The generalized **Toda** lattice

$$\psi(x+i\hbar) + \psi(x-i\hbar) + 2 \cosh x \psi(x) = E\psi(x)$$

- The situation is similar to the difference between the **Mathieu (cos)** and the **modified Mathieu (cosh)** potentials

To Calabi-Yau

$$E = e^{ik_x a} + e^{-ik_x a} + e^{ik_y a} + e^{-ik_y a}$$

$$E = e^p + e^{-p} + e^x + e^{-x}$$



$$X + X^{-1} + Y + Y^{-1} = E$$

**This equation defines a genus one
Riemann surface**

[Dunne's talk]

To Calabi-Yau

- **The complex 3d space** [Schiappa's talk]


$$VW = X + X^{-1} + Y + Y^{-1} - E$$

describes a Calabi-Yau manifold

- **The Riemann surface has enough information to describe this CY manifold**
- **In this way, one can see a connection to the CY geometry**

Mirror Symmetry

- The Calabi-Yau geometry has a remarkable hidden duality, called **mirror symmetry**



$$A : \text{Local } \mathbb{C}P^1 \times \mathbb{C}P^1$$

$$B : VW = X + X^{-1} + Y + Y^{-1} - E$$

Spectral Solution 1

- **The spectral problem of the N=2 generalized Toda lattice is solved by the exact version of the quantization condition in terms of string theory**

Grassi, YH & Marino '14; Wang, Zhang & Huang '15

$$\frac{\partial}{\partial t} F_{\mathbb{P}^1 \times \mathbb{P}^1}(t; \hbar) + \frac{\partial}{\partial \tilde{t}} F_{\mathbb{P}^1 \times \mathbb{P}^1}(\tilde{t}; \tilde{\hbar}) = 2\pi \left(n + \frac{1}{2} \right)$$

$$t = t(E; q) \quad q = e^{i\hbar}, \quad \tilde{t} = \frac{2\pi t}{\hbar}, \quad \tilde{\hbar} = \frac{4\pi^2}{\hbar}$$

Spectral Solution 2

- On the other hand, the spectrum of the Hofstadter problem is encoded in the **quantum deformed mirror map**


YH, Katsura & Tachikawa '16

$$\rho(E) = \frac{1}{2\pi} \operatorname{Im} \left[\frac{\partial t(E; q)}{\partial E} \right], \quad q = e^{2\pi i \alpha}$$

4. Semiclassical Analysis

@ Weak Coupling

- In the **weak magnetic regime**, the band width of the Hofstadter model is extremely narrow, and one can see Landau level splitting



@ Weak Coupling

- **This fact is easily seen by the Hamiltonian analysis**

$$\begin{aligned} H &= e^{\frac{ia}{\hbar}\Pi_x} + e^{-\frac{ia}{\hbar}\Pi_x} + e^{\frac{ia}{\hbar}\Pi_y} + e^{-\frac{ia}{\hbar}\Pi_y} \\ &= 4 - \frac{a^2}{\hbar^2} (\Pi_x^2 + \Pi_y^2) \\ &\quad + \frac{a^4}{12\hbar^4} (\Pi_x^4 + \Pi_y^4) + \dots \end{aligned}$$

@ Weak Coupling

- **This fact is easily seen by the Hamiltonian analysis**

$$\begin{aligned} H &= e^{\frac{ia}{\hbar}\Pi_x} + e^{-\frac{ia}{\hbar}\Pi_x} + e^{\frac{ia}{\hbar}\Pi_y} + e^{-\frac{ia}{\hbar}\Pi_y} \\ &= 4 - \frac{a^2}{\hbar^2} (\Pi_x^2 + \Pi_y^2) \\ &\quad + \frac{a^4}{12\hbar^4} (\Pi_x^4 + \Pi_y^4) + \dots \end{aligned}$$

$$E_n = 4 - (2n + 1)\phi + \frac{2n^2 + 2n + 1}{8}\phi^2 + \mathcal{O}(\phi^3)$$

$(\phi = 2\pi\alpha)$


@ Weak Coupling

- **This fact is easily seen by the Hamiltonian analysis**

$$\begin{aligned} H &= e^{\frac{ia}{\hbar}\Pi_x} + e^{-\frac{ia}{\hbar}\Pi_x} + e^{\frac{ia}{\hbar}\Pi_y} + e^{-\frac{ia}{\hbar}\Pi_y} \\ &= 4 - \frac{a^2}{\hbar^2} (\Pi_x^2 + \Pi_y^2) \\ &\quad + \frac{a^4}{12\hbar^4} (\Pi_x^4 + \Pi_y^4) + \dots \end{aligned}$$

$$E_n = 4 - (2n + 1)\phi + \frac{2n^2 + 2n + 1}{8}\phi^2 + \mathcal{O}(\phi^3)$$

$(\phi = 2\pi\alpha)$



@ Weak Coupling

- **There is a systematic way to compute the weak coupling expansion**

BenderWu package: Sulejmanpasic & Ünsal '16


Extension: Gu & Sulejmanpasic '17

$$E_0 = 4 - \phi + \frac{\phi^2}{8} - \frac{\phi^3}{192} + \frac{\phi^4}{768} + \frac{67\phi^5}{245760} + \frac{653\phi^6}{5898240} \\ + \dots + 6.79 \times 10^{39} \phi^{99} + 4.59 \times 10^{40} \phi^{100} + \mathcal{O}(\phi^{101})$$

- **Obviously, this expansion looks **divergent****
- **One needs a resummation method**


Borel(-Pade) Singularities

$$\mathcal{B}E_0(\zeta) := \sum_{n=0}^{\infty} \frac{E_0^{(n)}}{n!} \zeta^n$$



Borel(-Pade) Singularities

$$\mathcal{B}E_0(\zeta) := \sum_{n=0}^{\infty} \frac{E_0^{(n)}}{n!} \zeta^n$$




Borel(-Pade) Resum


$$\mathcal{S}_{\pm} E_0 := \frac{1}{\phi} \int_{C_{\pm}} d\zeta e^{-\zeta/\phi} \mathcal{B}E_0(\zeta)$$



$$\phi = \frac{2\pi}{5} \quad \left(\alpha = \frac{1}{5} \right)$$



$$\mathcal{S}_{\pm} E_0 = 2.935649214 \pm 0.000378867i$$




**This ambiguity should be canceled
by “nonperturbative” corrections**

What does this value mean?

Observation

YH, in progress

- The Borel resummed value is very close to the energy at the **Van Hove singularity**



Observation

- **For $\alpha = 1/5$, the positions of the Van Hove singularities are analytically determined by**

$$E \left(E^4 - 10E^2 + \frac{35 - 5\sqrt{5}}{2} \right) = 0$$

Observation

- **For $\alpha = 1/5$, the positions of the Van Hove singularities are analytically determined by**

$$E \left(E^4 - 10E^2 + \frac{35 - 5\sqrt{5}}{2} \right) = 0$$

$$E_{\text{VHS}} = \pm 2.935648819, \pm 1.175570505, 0$$

Observation

- **For $\alpha = 1/5$, the positions of the Van Hove singularities are analytically determined by**

$$E \left(E^4 - 10E^2 + \frac{35 - 5\sqrt{5}}{2} \right) = 0$$

$$E_{\text{VHS}} = \pm \underline{2.935648819}, \pm 1.175570505, 0$$

$$S_{\pm} E_0 = \underline{2.935649214} \pm 0.000378867i$$

Observation

- **The coincidence is probably not accidental**

$$\alpha = 1/10$$

$$E_{\text{VHS}}^{\text{max}} = \mathbf{3.41997695020118566}$$

$$S_{\pm}E_0 = \mathbf{3.41997695020118576} \pm 4.0 \times 10^{-9}i$$

$$\alpha = 1/15$$

$$E_{\text{VHS}}^{\text{max}} = \mathbf{3.602714983890327032980205}$$

$$S_{\pm}E_0 = \mathbf{3.602714983890327032980207} \pm 3.6 \times 10^{-14}i$$

Summary

- The **Hofstadter** model (2d electrons) and the generalized **Toda** lattice (integrable system) has a nontrivial relation to the **Calabi-Yau** geometry
- The weak magnetic expansion in the Hofstadter model is **not Borel summable**

Open Questions

- **The full weak magnetic expansion must be a **transseries** expansion**
- **P/NP (Dunne-Ünsal) relations?**
- **If α is **irrational**, the spectrum is much more involved**
- **What does the Borel resum for the irrational case mean?**

Open Questions

- **The full weak magnetic expansion must be a **transseries** expansion**
- **P/NP (Dunne-Ünsal) relations?**
- **If α is **irrational**, the spectrum is much more involved**
- **What does the Borel resum for the irrational case mean?**

I need your help!

Thank you!