

First Spectroscopy of ^{40}Mg

Monday, 4 June 2018 13:48 (18 minutes)

The study of nuclei far from stability is one of the most active and challenging areas of nuclear structure physics. One of the most exotic neutron-rich nuclei currently accessible to experiment is ^{40}Mg [1], which lies at the intersection of the nucleon magic number $N=28$ and the dripline, and is expected to have a large prolate deformation similar to that observed in the neighboring lighter isotopes $^{32-38}\text{Mg}$ [2]. In addition, the occupation of the weakly bound $p_{3/2}$ state may lead to the appearance of an extended neutron halo [3]. Thus ^{40}Mg offers an exciting possibility and a rare opportunity to investigate the coupling of weakly bound valence particles to a deformed core, and the influence of near threshold effects on collective rotational motion.

We will discuss the results of an experiment carried out at RIBF RIKEN to study low-lying states in ^{40}Mg produced by a 1-proton removal reaction from a ~ 240 MeV/u ^{41}Al secondary beam. ^{40}Mg and other final products were separated and identified using the Zero Degree Spectrometer, and prompt gamma rays detected using the DALI2 array. The observed excitation spectrum is shown to reveal unexpected properties as compared to both neighboring (more bound) Mg isotopes and theoretical model predictions.

This work is supported by the U.S. Department of Energy, Office of Nuclear Physics, under contract no DE-AC02-05CH11231.

[1] H. L. Crawford et al., Phys. Rev. C 89, 041303(R) (2014).

[2] P. Doornenbal et al., Phys. Rev. Lett. 111, 212502 (2013).

[3] F. Nowacki and A. Poves, Phys. Rev. C 79, 014310 (2009).

Summary

Primary authors: CRAWFORD, Heather (Lawrence Berkeley National Laboratory); Dr FALLON, Paul (Lawrence Berkeley Laboratory)

Presenter: CRAWFORD, Heather (Lawrence Berkeley National Laboratory)

Session Classification: Session 3