

In-beam gamma-ray spectroscopy of 51K and 53K

Friday, 8 June 2018 11:39 (18 minutes)

One of the major focus of modern nuclear physics is to explore which part of the nuclear interaction gives rise to significant shell modifications. Recently, the evolution of the $2s_{1/2}+$ and $1d_{3/2}+$ single-particle states in odd-A K isotopes attract particular interests. The energy gap between these two states decrease continuously when neutrons fill $f_{7/2}$ orbit. Inversion of the ordering of the $2s_{1/2}$ and $1d_{3/2}$ orbits has been observed in ^{47}K ($N=28$) and ^{49}K ($N=30$). As the neutrons continue filling the orbits beyond the $N = 28$ shell, reinversion was observed for the first time in ^{51}K using laser spectroscopy. Such reinversion is consistent with the shell model calculations using different effective interactions and was revealed to be mainly driven by the central term of the monopole interaction. However, different interactions predict very different energy gaps between $2s_{1/2}$ and $1d_{3/2}$ in ^{51}K . In addition, the shell model calculation and the recently available ab initio calculation also predict the reversion in ^{53}K but also with very different energy gaps. The experimental spectroscopy of the excited states in ^{51}K and ^{53}K are thus crucial to benchmark the shell model and ab initio calculations and improve our understanding on the shell evolution mechanisms.

The in-beam gamma-ray spectroscopy measurement of ^{51}K and ^{53}K was carried out at RIBF at RIKEN, as part of the third campaign of the SEASTAR program. The low-lying states of ^{51}K and ^{53}K were populated via $^{52}\text{Ca}(p, 2p)$ and $^{54}\text{Ca}(p, 2p)$, respectively. In the presentation, I will report on the energy level scheme of $^{51,53}\text{K}$, exclusive cross sections and the individual parallel momentum distributions.

Primary author: Dr SUN, Yelei (CEA Saclay)

Presenter: Dr SUN, Yelei (CEA Saclay)

Session Classification: Session 16