Contribution ID: 8

Type: Poster

Cryogenic spectroscopy of sp3 defect states in carbon nanotubes

Monday, 9 July 2018 17:00 (15 minutes)

Photoactive defect states in semiconducting single-walled carbon nanotubes (CNTs) have the potential to enable novel applications in quantum photonic technologies. While early experiments have established cryogenic CNTs as quantum light emitters [1], more recent work has identified luminescent defect states as sources of single photons up to room temperature [2,3]. For chemically-engineered CNTs, sp³ functionalization offers means to deterministically influence the corresponding trap state properties via the characteristics of the covalent side chains. Recent findings show that the incorporation of sp³ alkyl defects into the sp² lattice of narrow-diameter CNTs promotes the formation of both neutral and charged localized excitons [4] with unexpectedly high quantum efficiency. Here, we report the results of our cryogenic photoluminescence studies carried out on individual nanotubes functionalized with sp³ side-wall chemistry.

[1] A. Högele, C. Galland, M. Winger, A. Imamoglu, Phys. Rev. Lett. 100, 217401 (2008).

[2] X. Ma, N. F. Hartmann, J. K. S. Baldwin, S. K. Doorn, H. Htoon, Nat. Nanotech. 10, 671–675 (2015).

[3] X. He et al., Nat. Photon. 11, 577-582 (2017).

[4] H. Kwon, M. Kim, M. Nutz, N. F. Hartmann, V. Perrin, B. Meany, M. S. Hofmann, C. W. Clark, S. K. Doorn, A. Höegele, Y. Wang, submitted.

Summary

Primary author: NUTZ, Manuel (Ludwig-Maximilians-University Munich)

Co-authors: KWON, Hyejin (University of Maryland); KIM, Mijin (University of Maryland); WANG, YuHuang (University of Maryland); HÖGELE, Alexander (Fakultät für Physik, Munich Quantum Center, and Center for NanoScience (CeNS), Ludwig-Maximilians-Universität, Geschwister-Scholl-Platz 1, 80539 München, Germany)

Presenter: NUTZ, Manuel (Ludwig-Maximilians-University Munich)

Session Classification: Poster Session