

Cavity-control of bright and dark interlayer excitons in van der Waals heterostructures

Monday, 9 July 2018 17:45 (15 minutes)

Monolayer transition metal dichalcogenides integrated in optical microcavities host exciton-polaritons as a hallmark of the strong light-matter coupling regime [1]. Analogous concepts for hybrid light-matter systems employing spatially indirect excitons with a permanent electric dipole moment in heterobilayer crystals promise realizations of exciton-polariton gases and condensates with immanent dipolar interactions. Here, we identify optical signatures of spatially indirect momentum-bright and momentum-dark interlayer excitons in vertical MoSe₂-WSe₂ heterostructures and implement cavity-control of both exciton manifolds. To this end we employ a tunable open-access cavity with one curved fiber-based mirror [2] and one planar mirror with extended MoSe₂-WSe₂ heterobilayer flakes on top. The configuration of controlled intermirror spacing and lateral scanning capabilities is used to explore the light-matter coupling of excitons as a function of the cavity length at representative positions of heterobilayers selected by two-dimensional cavity imaging. Our experiments quantify the strength of interlayer excitons and demonstrate Purcell enhancement in cavity-modified photonic environments [3].

- [1] X. Liu, T. Galfsky, Z. Sun, F. Xia, E.-c. Lin, Y.-H. Lee, S. Kena-Cohen, and V. M. Menon, *Nature Photonics* 9, 30 (2015).
- [2] D. Hunger, T. Steinmetz, Y. Colombe, C. Deutsch, T. W. Hänsch, and J. Reichel, *New J. Phys.* 12,065038 (2010).
- [3] M. Förög, L. Colombier, R. Patel, J. Lindlau, A. D. Mohite, H. Yamaguchi, D. Hunger, and A. Högele, arXiv:1710.00990 (2017).

Summary

Primary author: Mr FÖRG, Michael (Ludwig-Maximilians-Universität Munich)

Co-authors: Mr COLOMBIER, Léo (Ludwig-Maximilians-Universität Munich); Mr PATEL, Robin K. (Ludwig-Maximilians-Universität Munich); Ms LINDLAU, Jessica (Ludwig-Maximilians-Universität Munich); Mr MOHITE, Aditya D. (Los Alamos National Laboratory); Mr YAMAGUCHI, Hisato (Los Alamos National Laboratory); Prof. HUNGER, David (Karlsruher Institut für Technologie); HÖGELE, Alexander (Fakultät für Physik, Munich Quantum Center, and Center for NanoScience (CeNS), Ludwig-Maximilians-Universität, Geschwister-Scholl-Platz 1, 80539 München, Germany)

Presenter: Mr FÖRG, Michael (Ludwig-Maximilians-Universität Munich)

Session Classification: Poster Session