

Studies of neutron-rich nuclear structures through beta-delayed decay of spin-polarized isotopes

Monday, 20 December 2010 13:30 (30 minutes)

Much attention has been paid on the exotic structure of neutron-rich nuclei such those around magic numbers $N = 8$ and 20 . However, most of the information on the excited states of these nuclei, such as spin and parity, has not been known well. We have developed a new method to effectively investigate the level structure by taking advantage of asymmetric beta-decay of spin polarized unstable nuclei: The spins of the daughter states can be assigned unambiguously from the characteristic asymmetry.

We have started beta-delayed decay spectroscopy at ISAC-1 TRIUMF, where highly polarized radioactive nuclear beams are available.

In the first experiment measuring beta-delayed neutron decays of spin-polarized ^{11}Li has successfully assigned the spins and parities of 7 levels in ^{11}Be for the first time [1].

The experiment with polarized ^{28}Na and ^{29}Na beams have been performed in 2007. The beta-decay asymmetry parameters and gamma-ray intensities have also assigned spin-parity of a newly found level in ^{28}Mg and of 7 levels in ^{29}Mg for the first time. The observed levels and log-ft values were compared with the shell model calculations using NuShell code with USD interactions. The level energies, log-ft values and the decay properties of all the assigned levels were explained well by assuming sd-shell configurations. However, in ^{29}Mg two levels at 1.095 and 1.430 MeV associated with large log-ft values could not be reproduced by the calculations. The Monte Carlo Shell Model calculation taking into account the intruder configurations predicted $3/2^-$ and $7/2^-$ levels around 1 MeV [2]. This fact strongly suggests negative parity assignments for the 1.095 and 1.430 MeV levels in ^{29}Mg . In August 2010 the experiment with ^{30}Na beam has been performed and the data analysis is in progress now.

In the talk the principle of the method will be introduced and results on the ^{11}Be , ^{28}Mg and ^{29}Mg structures will be discussed. Some of new findings on ^{30}Mg structure will also be presented.

[1] Y. Hirayama et al., Phys. Lett. B611, 239 (2005).

[2] Y. Utsuno, private communication.

Presenter: Prof. SHIMODA, Tadashi (Department of Physics, Osaka University)