

Shape coexistence studies in neutron-rich krypton isotopes around N=60

Friday, 12 April 2019 11:30 (15 minutes)

Even-even strontium and zirconium nuclei in the A=100 region show a sudden onset of deformation at N=60 while the lighter isotopes up to N=58 are rather spherical. Unlike, the even krypton isotopes exhibit a smooth onset of collectivity up to N=60 [1]. Recent high-resolution gamma-spectroscopy results on ^{96}Kr [2] following projectile fission confirmed the energy of the yrast $2^+ \rightarrow 0^+$ transition [1] and reported for the first time on the yrast $4^+ \rightarrow 2^+$ transition. Further on, recent results from the SEASTAR 2015 campaign on the extremely exotic $^{98,100}\text{Kr}$ isotopes, populated in nucleon knockout reactions of radioactive beams, reported a decrease in the energy of the yrast 2^+ states suggesting a continuation of the smooth shape transition beyond N=60 [3]. The side product data on the neutron-rich ^{96}Kr collected from the same campaign suggested new low-lying excited states [4], which unlike the oblate shape g.s., could correspond to the prolate minimum of the potential energy surface (see ref.[5] for example). Unfortunately the low-resolution data from DALI2 together with the possible lineshape effects due to state lifetimes, made it difficult to establish a reliable level scheme. We performed a subsequent experiment to excite the lowest non-yrast states of ^{96}Kr using Coulomb-nuclear excitation reactions at HIE-ISOLDE, CERN, which suffered from a reduced radioactive beam intensity and increased stable beam contamination.

As shown in the neighbouring nuclei of higher Z, the prolate shape is related to the lowering of the neutron $g_{9/2}$ orbital. Thus studies of the odd-neutron krypton isotopes $^{95,97}\text{Kr}$ are complementary and of similar importance. The existence of a $T_{1/2} = 1.4$ us isomer in ^{95}Kr [6] made it mandatory to measure delayed gamma-rays at the end of Zero Degree spectrometer, which we did in the SEASTAR 2015 campaign. The correlation of delayed and prompt gamma-rays worked nice, but again due to the low-resolution of DALI2 and the increased line density of the prompt spectrum only a preliminary level scheme could be built [7].

From the point of view of reaction mechanism and the production yields, the RIBF is currently the only place in the world where these shape-coexisting states in the krypton isotopes around N=60 could be studied.

Thus, in order to obtain definite results on the gamma-energies and their coincidence relations, as well as have a more sensitive analysis of the lineshapes, we propose to study the neutron-rich krypton isotopes around N=60 with high resolution gamma-ray spectroscopy at the RIBF populated via nucleon knockout reactions.

- [1] M. Albers et al., Phys. Rev. Lett. 108, 62701 (2012)
- [2] J. Dudouet et al., Phys. Rev. Lett. 118, 162501 (2017)
- [3] F. Flavigny et al., Phys. Rev. Lett. 118, 242501 (2017)
- [4] K. Moschner et al., in preparation.
- [5] K. Nomura et al., Phys. Rev. C 96, 034310 (2017)
- [6] J. Genevey et al., Phys. Rev. C 73, 37308 (2006)
- [7] R.-B. Gerst et al., in preparation.

Primary author: BLAZHEV, Andrey (Institute for Nuclear Physics, University of Cologne, Germany)

Presenter: BLAZHEV, Andrey (Institute for Nuclear Physics, University of Cologne, Germany)

Session Classification: Proposals