Contribution ID: 4

Study of unbound nuclei ³³Ne via one-proton knockout reactions

Friday, 30 August 2019 13:50 (20 minutes)

The magicity of N=20 in the vicinity of Ne, Na, and Mg isotopes vanishes due to *pf*-shell intruder configuration, which is called 'island of inversion' [1-3]. In recent years, shell evolution of Ne isotopes in this region is emerging topic of interest [4,5]. Nevertheless, there is no observed state of ³³Ne.

It is only known that ³³Ne is unbound nuclei [6] and the *1n* separation energy S_n is only predicted to -0.9 MeV [7].

The experiment was carried out at the RIBF in RIKEN. The secondary beam of ³⁴Na at 264 MeV/nucleon was provided by BigRIPS [8] and impinged on the carbon reaction target. After the one-proton knockout reaction of ³⁴Na, ³³Ne was produced and immediately decayed into ³²Ne and a neutron. The invariant mass spectrum of ³²Ne + *n* system was reconstructed by measurement of fragments and neutrons using SAMURAI spectrometer [9]. In this presentation, details of analysis and preliminary results of ³²Ne + *n* invariant mass spectrum will be discussed.

- [1] Z. Elekes et al., Phys. Rev. C 73, 044314 (2006).
- [2] P. Doornenbal et al., Phys. Rev. Lett. 103, 032501 (2009).
- [3] P. Doornenbal et al., Phys. Rev. C 81, 041305 (2010).
- [4] T. Nakamura et al., Phys. Rev. Lett. 112, 142501 (2014).
- [5] I. Murray et al., Phys. Rev. C 99, 011302 (2019).
- [6] M. Notani et al., Phys. Lett. B 542, 49 (2002).
- [7] G. Audi, H. Wapstra, and C. Thibault, Nucl. Phys. A 729, 337 (2003).
- [8] T. Kubo, Nucl. Inst. and Meth. B 204, 97 (2003).
- [9] T. Kobayashi et al., Nucl. Inst. and Meth. B 317, 294 (2013).

Primary author: Mr CHAE, Hyunwoo (Seoul National University)

Presenter: Mr CHAE, Hyunwoo (Seoul National University)

Session Classification: Status report