

Contribution ID: 92

Type: **Parallel Session Presentation**

Transverse single-spin asymmetries of W_{\pm} -bosons in p+p collisions at $\sqrt{s} = 510$ GeV

Friday, 22 October 2021 08:12 (18 minutes)

The STAR experiment at RHIC has measured transverse single-spin asymmetries of W_{\pm} -bosons in proton-proton collisions at a center-of-mass energy $\sqrt{s} = 510$ GeV (2017 data). These asymmetries probe correlations between parton motion and the proton spin in the initial state which are described in terms of transverse momentum dependent parton distribution functions (TMD), in this case the Sivers function. The Sivers function is of particular theoretical interest because its process dependence can be linked to underlying kinematics, namely the gauge link structure of the scattered parton with the nucleon remnant. This means that the Sivers function is not universal and a sign change is expected between the asymmetries measured in semi-inclusive deep inelastic scattering compared to those in hadronic collisions. The new STAR preliminary results with an integrated luminosity of about 350 pb^{-1} improve significantly on previous data from 2011. We will discuss details of the full reconstruction of the W -boson kinematics which are required for a true TMD measurement. Comparison with recent global fits will illustrate the potential impact of the new data.

Primary authors: EYSER, K. Oleg (UC Riverside); STAR COLLABORATION

Presenter: EYSER, K. Oleg (UC Riverside)

Session Classification: Transverse Momentum Structure (TMD)

Track Classification: Parallel Sessions: Transverse momentum structure (TMD)