

Roles of spin-dependent transitions in nuclei on astrophysical processes in stars

Wednesday, 20 October 2021 12:25 (20 minutes)

Due to recent advances in shell-model studies on spin modes in nuclei, precise evaluations of Gamow-Teller (GT) strengths become feasible and electron-capture and β -decay rates in stellar environments have been updated.

The weak rates in *sd*-shell obtained with the USDB Hamiltonian are applied to nuclear Urca processes in O-Ne-Mg cores in stars with 8-10 solar masses [1,2]. The Urca processes for the nuclear pairs with $A=23$ and 25 are found to be important for the cooling of the core [1].

Here, the weak rates important for the Urca processes in accreted neutron star crusts [3] are investigated. The e-capture and β -decay rates are evaluated for the nuclear pair with $A=31$, $^{31}\text{Al} \leftrightarrow ^{31}\text{Mg}$, in *sd-pf* shell and the pair with $A=61$, $^{61}\text{V} \leftrightarrow ^{61}\text{Cr}$, in *fp-gd* shell. ^{31}Mg belongs to the island of inversion, where admixtures of *sd*- and *fp*-shells become important. Energy levels in ^{31}Mg are found to be well reproduced with the use of EEdf1 interaction obtained by the extended Kuo-Krensiglowa (EKK) method [4], which can properly treat Q-box calculations in two-major shells without divergence problems. The weak rates evaluated with the EKK method prove to lead to Urca processes.

The GT strengths in ^{61}V is evaluated with the GXPF1J Hamiltonian [5]. The calculated strength between the ground states of ^{61}V and ^{61}Cr is found to be consistent with the recent experimental data [6]. This suggests that the Urca process for the $A=61$ pair would be more moderate than considered before. Results with an extension to the *fp-gd* shell-model space will be also reported.

The weak rates in *fp*-shell obtained with the GXPF1J are applied to nucleosynthesis in Type Ia supernova explosions [7]. The electron screening effects are taken into account [8]. Overproduction problem of neutron-rich iron-group elements for the previous weak rates is found to be considerably suppressed.

[1] H. Toki, T. Suzuki, K. Nomoto, S. Jones, and R. Hirschi, Phys. Rev. C **88**, 015806 (2013).

[2] T. Suzuki, H. Toki, and K. Nomoto, ApJ. **817**, 163 (2016).

[3] H. Schatz, S. Gupta, P. Moller et al., Nature **50**, 62 (2014).

[4] N. Tsunoda, T. Otsuka, N. Shimizu, M. Hjorth-Jensen, K. Takayanagi, and T. Suzuki, Phys. Rev. C **95**, 021304 (R) (2017).

[5] M. Honma et al., J. Phys. Conf. Ser. **20**, 7 (2005).

[6] W. J. Ong et al., Phys. Rev. Lett. **125**, 262701 (2020).

[7] K. Mori et al., M. A. Famiano, T. Kajino et al., ApJ. **863**, 176 (2018);

[8] K. Mori, T. Suzuki, M. Honma et al., ApJ **904**, 29 (2020).

Primary authors: Prof. SUZUKI, Toshio (Nihon University); Prof. HONMA, Michio (Aizu University); Prof. SHIMIZU, Noritaka (CNS, The University of Tokyo); Prof. OTSUKA, Takaharu (The University of Tokyo, and RIKEN); Dr MORI, Kanji (Fukuoka University); Prof. KAJINO, Toshitaka (Beihan University)

Presenter: Prof. SUZUKI, Toshio (Nihon University)

Session Classification: Spin in Nuclear Reactions and Nuclei

Track Classification: Parallel Sessions: Spin in Nuclear Reactions and Nuclei