

Symposium on Nuclear Data 2020

Ag102 12.9 m	Ag103 65.7 m	Ag104 69.2m	Ag105 41.29 d	S ymposium on	Ag107 51.839 %	Ag108 2.37 m	Ag109 48.161 %	Ag110 24.6 s	Ag111 7.45 d	Ag112 2.135 h
Pd101 8.47 h	Pd102 1.02 %	Pd103 16.991 d	Pd104 11.14 %	Pd105 22.33 %	N uclear	Pd107 6.36×4 y	Pd108 26.46 %	Pd109 15.700(26)	Pd110 11.72 %	Pd111 33.4 m
Rh100 20.8 h	Rh101 3.3 y	Rh102 2.72 d	Rh103 100 %	Rh104 42.3 s	Rh105 35.95 h	D ata	2020 Nov.	Rh108 9.0 m	Rh109 99 s	Rh110 3.3 s

Contribution ID: 39

Type: **Poster Presentation**

Estimation of Flux and Residual Radioactivity for the COMET Phase-I Experiment / COMET Phase-I 実験のための粒子線束と残留放射能の評価

Thursday, 26 November 2020 17:11 (1h 39m)

COherent Muon to Electron Transition (COMET) is an experiment at J-PARC, which will search for coherent neutrino-less conversion of a muon to an electron in muonic atoms. The experiment will be carried out in two steps: Phase-I and Phase-II. In the Phase-I experiment, 3.2 kW 8 GeV proton beam irradiates a 70-cm long graphite target to produce negative pions. The negative pions are captured in the magnetic field and delivered to pion-decay and muon-transport sections. The Phase-I experiment aims to detect the $\mu^- e$ conversion events and measure the beam-related background events for the Phase-II experiment. Now, it has been planned that the maintenance by radiation workers would be conducted after the 150-day operation and the following 180-day cooling. It is necessary to evaluate the residual radiation dose for the safety of the workers during the maintenance. On this study, we calculated fluxes of neutron, photon, proton and other charged particles in the beam room during the beam operation and the residual activity after the cooling time by using Monte Carlo simulation code PHITS version 3.22 and DCHAIN-PHITS version 3.21. The calculation results show that the design of components around the target and beam dump needs to be improved to reduce the radioactivities after the cooling time.

Primary author: TOKUNAGA / 徳永, Naoki / 直輝 (Kyushu University / 九州大学)

Co-authors: SHIGYO / 執行, Nobuhiro / 信寛 (Kyushu University / 九州大学); MAKIMURA / 牧村, Shunsuke / 俊助 (Institute of Particle and Nuclear Studies J-PARC Center Particle and Nuclear Physics Division Hadron Section Tokai campus / 素粒子原子核研究所 J-PARC センター素粒子原子核ディビジョンハドロンセクション東海キャンパス); FUKAO / 深尾, Yoshinori / 祥紀 (Institute of Particle and Nuclear Studies J-PARC Center Particle and Nuclear Physics Division Hadron Section Tokai campus / 素粒子原子核研究所 J-PARC センター素粒子原子核ディビジョンハドロンセクション東海キャンパス); KAMEI / 亀井, Naoya / 直矢 (Institute of Particle and Nuclear Studies J-PARC Center Particle and Nuclear Physics Division Hadron Section Tokai campus / 素粒子原子核研究所 J-PARC センター素粒子原子核ディビジョンハドロンセクション東海キャンパス); YOSHIDA / 吉田, Makoto / 誠 (J-PARC Center, High Energy Accelerator Research Organization (KEK) / 高エネルギー加速器研究機構 J-PARC センター)

Presenter: TOKUNAGA / 徳永, Naoki / 直輝 (Kyushu University / 九州大学)

Session Classification: Poster