

## Symposium on Nuclear Data 2020

Ag102 12.9 m	Ag103 65.7 m	Ag104 69.2m	Ag105 41.29 d	<b>S</b> ymposium on	Ag107 51.839 %	Ag108 2.37 m	Ag109 48.161 %	Ag110 24.6 s	Ag111 7.45 d	Ag112 2.195 h
Pd101 8.47 h	Pd102 1.02 %	Pd103 16.991 d	Pd104 11.14 %	Pd105 22.33 %	<b>N</b> uclear	Pd107 6.5e+4 y	Pd108 26.46 %	Pd109 15.700(26)	Pd110 11.72 %	Pd111 23.4 m
Rh100 20.8 h	Rh101 3.3 y	Rh102 2.72 y	Rh103 100 %	Rh104 42.3 s	Rh105 35.98 h	<b>D</b> ata	<b>2020</b> Nov.	Rh108 9.0 m	Rh109 99 s	Rh110 3.3 s

Contribution ID: 52

Type: **Poster Presentation**

# Nuclide production cross sections of natLu target irradiated with 0.4-, 1.3-, 2.2-, 3.0-GeV protons/natLu 標的に対する 0.4, 1.3, 2.2, 3.0 GeV 陽子入射による核種生成断面積

Thursday, 26 November 2020 17:10 (1h 40m)

Reliable assessment of radioactivity in target and structural materials for high-energy accelerator facilities such as accelerator-driven systems and spallation neutron sources requires detailed information on nuclide production cross sections by spallation reactions. To obtain the systematic cross section data for nuclide productions by spallation reactions, we have conducted irradiation experiments at Japan Proton Accelerator Research Complex (J-PARC). So far, we have measured nuclide production cross sections for light to medium-heavy target elements ( $Z \leq 47$ ) with proton energies ranging from 0.4 to 3.0 GeV. To investigate heavier target elements, we conducted an experiment for target elements with atomic number around  $Z=70$ , including  $^{nat}\text{Lu}$  ( $Z=71$ ) target.

Four sets of Ho (90mg/cm<sup>2</sup>), Lu (100mg/cm<sup>2</sup>), and Re (210mg/cm<sup>2</sup>) foils were packed in aluminum containers together with 0.1-mm-thick aluminum catchers to avoid recoil contamination. Each set of targets was irradiated with 0.4-, 1.3-, 2.2-, and 3.0-GeV protons accelerated by 3-GeV Rapid Cycling Synchrotron (RCS). The beam current was monitored by a current transformer installed in front of the irradiation position. After the irradiation, gamma-rays emitted from the samples were detected by two high-purity Germanium detectors (relative efficiency 20%, Canberra Co., Ltd.).

The measured cross sections were compared with theoretical predictions by Particle and Heavy Ion Transport code System (PHITS) [1], and INCL++/ABLA[2,3]. The figure shows experimental and calculated  $^{nat}\text{Lu}(p, X)^{nat}\text{Be}$  reaction cross sections. INCL/GEM model implemented in PHITS underestimated the experimental cross sections by a factor of about 2.

In the presentation, we will report our experimental results for the natLu target, and more detailed discussion on reaction mechanisms will be given.

## References

- [1] T. Sato, et al., J. Nucl. Sci. Technol. 55, pp.684 (2018).
- [2] D. Mancusi et al., Phys. Rev. C 91:034602 (2015).
- [3] A. Kelic et al., Proceedings of Joint ICTP-IAEA Advanced Workshop on Model Codes for Spallation Reactions, ICTP Trieste, Italy pp.181 (2008).

**Primary authors:** Mr TAKESHITA/竹下, Hayato/隼人 (Kyushu University); MEIGO, Shin-ichiro (J-PARC/JAEA); Dr MATSUDA, Hiroki (J-PARC Center, JAEA); IWAMOTO, Hiroki (Japan Atomic Energy Agency); NAKANO, Keita (JAEA); Prof. WATANABE, Yukinobu (Kyushu University); Dr MAEKAWA / 前川, Fujio / 藤夫 (JAEA / JAEA)

**Presenter:** Mr TAKESHITA/竹下, Hayato/隼人 (Kyushu University)

**Session Classification:** Poster