

Symposium on Nuclear Data 2020

Ag102 12.9 m	Ag103 65.7 m	Ag104 69.2m	Ag105 41.29 d	S ymposium on	Ag107 51.839 %	Ag108 2.37 m	Ag109 48.161 %	Ag110 24.6 s	Ag111 7.45 d	Ag112 3.130 h
Pd101 8.47 h	Pd102 1.02 %	Pd103 16.991 d	Pd104 11.14 %	Pd105 22.33 %	N uclear	Pd107 8.36 s	Pd108 26.46 %	Pd109 11.7002h	Pd110 11.72 %	Pd111 33.4 m
Rh100 20.8 h	Rh101 3.3 y	Rh102 2.72 h	Rh103 100 %	Rh104 42.3 s	Rh105 37.95 h	D ata	2020 Nov.	Rh108 3.0 m	Rh109 89 s	Rh110 3.3 s

Contribution ID: 36

Type: **Poster Presentation**

The origin of correlation between mass and angle in quasi-fission/準核分裂における質量と角度の相関の起源

Thursday, 26 November 2020 16:50 (1h 58m)

Mass-angle distribution (MAD) measurement of heavy and superheavy element fragmentation reactions is one of the powerful tools for investigating the mechanism of fission and fusion process. MAD shows a strong correlation between mass and angle when the quasi-fission event is dominant. It has characteristic that appears diagonal correlation as long as the quasi-fission event is dominant. This diagonal correlation could not be reproduced in previous our model before the introduction of the parameters.

In this study, we systematically evaluate the uncertainty model parameters contained in our model and clarify those model parameters to reproduce the diagonal correlation that appears in MAD. Using a dynamical model based on the fluctuation diffraction theorem that employs Langevin equations, we calculate the mass angle distribution and mass distribution of the four reaction systems $48\text{Ti} + 186\text{W}$, $34\text{S} + 232\text{Th}$, $48\text{Ti} + 208\text{Pb}$, and $28\text{S} + 238\text{U}$, which are dominated by quasi-fission. We were able to clarify the effects of uncertain model parameters on the mass angle distribution and mass distribution. In addition, we identified the values of model parameters that can reproduce the correlation between mass and angle. As a result, it found that the balance of tangential friction and moment of inertia values is important for the correlation between mass and angle.

Primary author: AMANO/天野, Shota/翔太 (Kindai University/近畿大学)

Co-authors: ARITOMO/有友, Yoshihiro/嘉浩 (Kindai University/近畿大学); ISHIZAKI/石崎, Shoma/翔馬 (Kindai university/近畿大学); OKUBAYASHI/奥林, Mizuki/瑞貴 (Kindai University/近畿大学); OKUGAWA/奥川, Seiya/誠也 (Kindai university/近畿大学)

Presenter: AMANO/天野, Shota/翔太 (Kindai University/近畿大学)

Session Classification: Poster