

Weak-binding relation in the zero range limit

Tuesday, 9 March 2021 15:10 (20 minutes)

The ordinary hadrons consist of two quarks for mesons or three quarks for baryons. The hadrons with the other structures such as the multi-quark states or the hadronic molecular states are called exotic hadrons. Here, we would like to determine the structure of exotic hadrons using the weak-binding relation for the near-threshold hadrons. In the weak-binding relation, we can determine compositeness of the hadron X defined as the weight of the hadronic molecule component. The relation is expressed by the observables such as the scattering length a_0 and the radius R . Determining X from the observables, we can analyze the internal structure of exotic hadrons without any model calculations. The relation has the correction terms, which are estimated by the typical length of the interaction R_{typ} . Therefore the correction terms vanish in the exact zero range limit $R_{\text{typ}}=0$. We show that there are exceptional cases which violate the weak-binding relation, when the zero range limit is taken with a fine tuning of parameters. We propose a suitable modification of the correction terms of the weak-binding relation with the redefinition of R_{typ} , so that the relation is valid also in the zero range limit. Originally, R_{typ} was given by the interaction range R_{int} , but we define R_{typ} as the largest value among R_{int} and R_{eff} . R_{eff} is defined as the length scale in the effective-range expansion expect for the scattering length a_0 . We apply the modified weak-binding relation to the concrete model and evaluate the correction terms. We expect that the the system with large R_{eff} such as $\Lambda_c(2595)$ can be analyzed with the modified weak-binding relation.

Presenter: KINUGAWA, Tomona (Department of Physics, Tokyo Metropolitan University)